

PHARMACEUTICAL USES AND SYNTHESIS OF BENZOBICYCLOOCTANES

This application claims the benefit of U.S. Provisional Patent Application No. 60/257,532, filed December 22, 2000, where this provisional application is
5 incorporated herein by reference in its entirety.

TECHNICAL FIELD

The present invention is generally directed to benzobicyclooctanes, their use in inhibiting cellular events involving TNF- α or IL-8, and in the treatment of inflammation events in general; the application also provides a combinatorial library of diverse
10 bicyclooctanes and methods for their synthesis in a library format as well as individual compounds.

BACKGROUND OF THE INVENTION

One process for discovering new therapeutically active compounds for a given indication involves the screening of all compounds from available compound
15 collections. From the compounds tested, one or more structures is selected as a promising lead. A large number of related analogues are then synthesized in order to develop a structure-activity relationship (SAR). The SARs direct the development and then selection of one or more optimal compounds following traditional one-at-a-time synthesis and biological testing. This optimization process is long and labor intensive.

20 Adding significant numbers of new structures to the compound collections used in this initial screening step of the discovery and optimization process cannot be accomplished with traditional one-at-a-time synthesis methods, except over a time frame of months or even years. Faster methods are needed that allow for the preparation of libraries of related compounds in a matter of days or a few weeks. This need is particularly
25 apparent when it comes to synthesizing more complex compounds.

Combinatorial approaches have recently been extended to "organic" or non-peptide, libraries. The organic libraries at present, however, are limited in diversity and

generally relate to peptidomimetic compounds; in other words, organic molecules that repeat a peptide chain pharmacophore. There is a need in the art for additional approaches to the preparation of new organic libraries.

Cytokines are pleiotropic extracellular proteins that are released and
5 recognized by a wide variety of cell types. Via a series of complex interactions they are responsible for regulating many of the events involved in growth and differentiation of an organism. Among the cytokines, tumor necrosis factor- α (TNF- α) has been shown to play an important role in the genesis of certain chronic inflammatory and autoimmune diseases. TNF- α is secreted mainly by macrophages and monocytes in response to a variety of
10 inflammatory agents. Other cell types such as NK cells, T cells, B cells, Kupfer cells, and glial cells also produce TNF- α .

TNF- α is synthesized as an inactive 26 kDa pro-protein which is cleaved by the metalloprotease TNF- α Converting Enzyme (TACE) to afford the active 17 kDa cytokine protein. The cytokine exerts its biological effects by interacting with two high
15 affinity receptors of molecular weights 55 kDa (TNFR1 or p55) and 75 kDa (TNFR2 or p75) found on the surface of most cell types. As a result of TNF- α binding to its receptors, a cascade of signaling events occurs within the cell. The exact nature and sequence of events is dependent upon cell type and receptor. Two of the most important physiological effects of TNF- α binding to its receptors are the upregulation of new genes by activation of
20 the transcription factor NF κ B, and induction of programmed cell death or apoptosis.

Apoptosis is a normal physiological process whereby incompetent cells become targeted for disposal by the immune system. The process involves a series of morphological changes to the apoptotic cell, including a change of surface chemistry. This change in surface chemistry is recognized by macrophages that rapidly phagocytose the
25 cell. A number of stimuli can induce apoptosis, including DNA damage, UV radiation, growth factor deprivation, bacterial and viral infection, and ligation of cell surface receptors. TNF- α has been shown to induce apoptosis by binding to TNFR1. Under normal biochemical circumstances the process of apoptosis is integral in regulating the homeostatic balance between cell death and cell proliferation. However in many

autoimmune diseases this balance is shifted; not only do unwanted cells undergo apoptosis but healthy cells as well. These diseases are often associated with increased levels of TNF- α . There is a need in the art for compounds that can modulate binding of TNF- α to cell receptors, and/or modulate the consequential intracellular events.

5 Transcription factors are a family of proteins that bind to DNA and serve to upregulate gene expression. Often they remain in an inactive form until acted upon by a biochemical signal. One such transcription factor is nuclear factor kappa B (NF κ B), which can be activated by the binding of TNF- α to TNFR1 and/or TNFR2. NF κ B regulates many of the cytokines and other proinflammatory molecules associated with inflammatory and
10 autoimmune diseases. Classes of proteins subject to regulation by NF κ B and which have been demonstrated to be involved with disease states are cytokines and growth factors, adhesion molecules, chemokines, and immunoreceptors.

The inhibition of TNF- α induced apoptosis and of NF κ B activation is one means of preventing and/or treating autoimmune and inflammatory diseases including, but
15 not limited to, rheumatoid arthritis, inflammatory bowel disease, psoriasis, atherosclerosis, asthma, reperfusion injury, ischemia, sepsis, graft vs. host disease, adult respiratory distress syndrome, multiple sclerosis, and a host of severe invasive infections such as fulminant hepatitis, AIDS and bacterial meningitis, and allergic inflammation of the lungs and airways.

Interleukin-8 (IL-8) is a chemokine (chemotactic cytokine) which plays an
20 important role in the recruitment of neutrophils to sites of inflammation. It is a member of the CXC subfamily of chemokines, members of which contain a single amino acid residue between the first two cysteines. In addition to inducing the chemotaxis of neutrophils, IL-8 exerts other immunomodulatory effects such as release of superoxide, mobilization of intracellular Ca⁺⁺, upregulation of cell surface integrins, and activation of phospholipase
25 D. All of these cellular events are the result of IL-8 binding to one of its two high affinity receptors. The two receptors, known as IL8RA or CXCR1 and IL8RB or CXCR2, bind the ligand with a K_d of ca. 2 nM.

Numerous reports in the literature have associated increased levels of IL-8 with the development of inflammatory and autoimmune diseases such as Inflammatory

Bowel Disease (IBD), psoriasis, rheumatoid arthritis, Acute Respiratory Distress Syndrome (ARDS), cancer, atherosclerosis, reperfusion injury, and graft vs. host disease. The inhibition of IL-8 or other CXC chemokines from binding to CXCR1 and/or CXCR2 receptors is one means of preventing and/or treating these diseases.

5 Although treatment regimens are available for the symptomatic amelioration of some or all of these diseases, there still exists the need for compositions and methods for preventing and/or treating the inflammation which is often associated with the disease.


 The present invention satisfies these needs and provides related advantages as well, as described more fully herein.

10 SUMMARY OF THE INVENTION

 The present invention overcomes the known limitations to classical organic synthesis of bicyclooctanes, and the shortcomings in applying combinatorial chemistry to bicyclooctanes, as well as providing compounds which are useful in inhibiting TNF- α , IL-8, apoptotic-mediated processes, and inflammatory conditions. Moreover, the present
15 invention provides a library of diverse bicyclooctanes useful in elucidating the structure-function relationship in biological processes, such as inflammation.

 In one embodiment, the present invention provides a compound of formula

(I)



(I)

20 and optical isomers, diastereomers, enantiomers and pharmaceutically acceptable salts thereof in isolation or mixture, where, independently at each location:

 R¹ is selected from the following six formulae:

R^2 is $-OR^9$ or $-NR^9R^9$;

- R^3 is selected from hydrogen, halogen, hydroxyl or protected hydroxyl, amino or protected amino, and C_1 - C_8 alkyl or C_1 - C_8 haloalkyl;

R^4 and R^5 are independently selected from R^9 , $-OR^9$, $-NR^9R^9$ and $-N=N-R^9$; or R^4 and R^5 may together form a group selected from $=O$, $=CR^8R^8$ and $=NR^{10}$, or R^4 and R^5 may together with the carbon to which they are both attached form a spiro carbocyclic or heterocyclic ring;

- R^6 is selected from hydrogen, inorganic groups having 1-8 atoms exclusively selected from boron, sulfur, phosphorous, silicon and hydrogen, and organic groups having 1-20 carbons and optionally containing 1-4 heteroatoms selected from nitrogen, oxygen and silicon;

- R^7 is selected from halogen, hydroxyl or protected hydroxyl, amino or protected amino, and C_1 - C_8 alkyl or C_1 - C_8 haloalkyl;

R^8 is selected from hydrogen, alkyl, aryl and heteroalkyl;


- R^9 is selected from hydrogen and organic groups having 1-30 carbons and optionally containing 1-4 heteroatoms selected from nitrogen, oxygen and silicon, with the provision that two R^9 groups both joined to a common atom may be joined together so as to form a ring with the common atom;

R^{10} is selected from $-R^9$, $-OR^9$, $-NR^9R^9$, $-NH-C(O)R^9$, $-NH-C(O)OR^9$ and $-NH-C(S)NHR^9$; and

n is 0, 1, 2 or 3;


with the proviso that when R^6 is hydrogen and R^4 and R^5 together form $=O$ and R^1 is CO_2R^9 , then R^2 is not OCH_3 .

In one embodiment R^1 is




5

In one embodiment R^1 is




In one embodiment R^1 is




In any of the above embodiments, R^8 is, in one embodiment, selected from hydrogen and C_1 - C_8 alkyl. In a further embodiment, R^8 is hydrogen.

In one embodiment R^1 is




10

In one embodiment R^1 is



In one embodiment R^1 is



In one embodiment, R^1 is selected from the following five formulae:

In one embodiment R^1 is selected from the following four formulae:

- In any of the above embodiments, in a further embodiment, R^9 is independently selected at each occurrence from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5, with the provision that

two R^9 groups both joined to a common atom may be joined together so as to form a ring with the common atom.

In any of the above embodiments, in a further embodiment, R^9 is independently selected at each occurrence from R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene, R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5.

In any of the above embodiments, in a further embodiment, R^9 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, (heteroaryl) C_1 - C_{15} alkylene, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, C_6 - C_{10} aryl fused to C_1 - C_{15} alkylene, (alkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, $(C_6$ - C_{10} aryl)(C_6 - C_{10} arylene) C_1 - C_{15} alkylene, $(C_1$ - C_{15} alkyl) $_p$ (heteroarylene) C_1 - C_{15} alkylene, and (heteroalkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, or two R^9 groups bonded to a common nitrogen of R^1 may be joined together to form a 5-8 membered heterocycle including the common nitrogen, where this 5-8 membered heterocycle may be substituted with 0-5 groups selected from alkyl and heteroalkyl, where p is selected from 1, 2, 3, 4 and 5.

In any of the above embodiments, in a further embodiment, R^9 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, (heteroaryl) C_1 - C_{15} alkylene, and (heteroalkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, or the two R^9 groups joined to a common nitrogen of R^1 may be joined together to form a 5-8 membered heterocycle including the common nitrogen.

In any of the above embodiments, in a further embodiment, R^9 is selected from heteroalkyl, C_1 - C_{15} alkyl, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, $(C_6$ - C_{10} aryl)(C_6 - C_{10} arylene) C_1 - C_{15} alkylene, $(C_1$ - C_{15} alkyl) $_p$ (heteroarylene) C_1 - C_{15} alkylene, and C_6 - C_{10} aryl fused to C_1 - C_{15} alkylene.

In any of the above embodiments, in a further embodiment, R^9 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, $(C_6$ - C_{10} aryl)(C_6 -

C₁₀arylene)C₁-C₁₅alkylene, (C₁-C₁₅alkyl)_p(heteroarylene)C₁-C₁₅alkylene, and C₆-C₁₀aryl fused to C₁-C₁₅alkylene.

In any of the above embodiments, in a further embodiment, R⁹ is selected from hydrogen, heteroalkyl, C₁-C₁₅alkyl, (heteroaryl)C₁-C₁₅alkylene, and (heteroalkyl)_p(C₆-C₁₀arylene)C₁-C₁₅alkylene.

In any of the above embodiments, in a further embodiment, R⁹ is selected from hydrogen, heteroalkyl, C₁-C₁₅alkyl, (heteroaryl)C₁-C₁₅alkylene, (C₆-C₁₀aryl)C₁-C₁₅alkylene, (alkyl)_p(C₆-C₁₀arylene)C₁-C₁₅alkylene, or the two R⁹ groups of R¹ may be joined together to form a 5-8 membered heterocycle including the common nitrogen, where this 5-8 membered heterocycle may be substituted with 0-5 groups selected from alkyl and heteroalkyl.

In any of the above embodiments, in a further embodiment, R² is -OR⁹.

In any of the above embodiments, in a further embodiment, R² is -NR⁹R⁹.

In any of the above embodiments, in a further embodiment, R⁹ of R² is selected from hydrogen, R¹¹, R¹², R¹³, R¹⁴ and R¹⁵ where R¹¹ is selected from alkyl, heteroalkyl, aryl and heteroaryl; R¹² is selected from (R¹¹)_p-alkylene, (R¹¹)_p-heteroalkylene, (R¹¹)_p-arylene and (R¹¹)_p-heteroarylene; R¹³ is selected from (R¹²)_p-alkylene, (R¹²)_p-heteroalkylene, (R¹²)_p-arylene, and (R¹²)_p-heteroarylene; R¹⁴ is selected from (R¹³)_p-alkylene, (R¹³)_p-heteroalkylene, (R¹³)_p-arylene, and (R¹³)_p-heteroarylene, R¹⁵ is selected from (R¹⁴)_p-alkylene, (R¹⁴)_p-heteroalkylene, (R¹⁴)_p-arylene, and (R¹⁴)_p-heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5.

In any of the above embodiments, in a further embodiment, R⁹ of R² is selected from hydrogen, heteroalkyl, C₁-C₁₅alkyl, (C₆-C₁₀aryl)(C₆-C₁₀arylene)C₁-C₁₅alkylene, (C₁-C₁₅alkyl)_p(heteroarylene)C₁-C₁₅alkylene, (C₁-C₁₅alkyl)_p(heteroarylene)-heteroalkylene, (heteroalkyl)_p(C₆-C₁₀arylene)C₁-C₁₅alkylene, and (C₁-C₁₅alkyl)_p(C₆-C₁₀arylene)heteroalkylene.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R² is -OR⁹ where R⁹ is selected from a

heteroalkyl group having preferably 1-10 carbons and 1-4 heteroatoms selected from nitrogen, oxygen, silicon and sulfur.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^2 is $-NR^9R^9$ and R^9 is selected from
5 hydrogen, heteroalkyl, C_1 - C_{15} alkyl, (heteroaryl) C_1 - C_{15} alkylene, (heteroalkyl) $_p$ (aryl)-heteroalkylene, (heteroalkyl) $_p$ (aryl) C_1 - C_{15} alkylene, and $(C_1$ - C_{15} alkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene.

In any of the above embodiments, in a further embodiment, R^3 is selected from hydrogen and alkyl.

10 In any of the above embodiments, in a further embodiment, R^3 is hydrogen.

In any of the above embodiments, in a further embodiment, R^4 and R^5 are independently selected from R^9 , $-OR^9$, $-NR^9R^9$ and $-N=N-R^9$.

In any of the above embodiments, in a further embodiment, R^9 of R^4 and R^5 is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl,
15 heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -
20 heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5.

In any of the above embodiments, in a further embodiment, each of R^4 and R^5 is hydrogen.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment, at least one of R^4 and R^5 is selected from C_1 -
25 C_{15} alkyl, heteroalkyl, and C_6 - C_{10} aryl.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment, one of R^4 and R^5 is hydrogen and the other of R^4 and R^5 is selected from hydrogen, $-OR^9$, $-NR^9R^9$ and $-N=N-R^9$ where the R^9 is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl

and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene, R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment, R^4 and R^5 together form a group selected from $=O$, $=CR^8R^8$ and $=NR^{10}$.

10 In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^4 and R^5 together form $=O$.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^4 and R^5 together form $=NR^{10}$ and R^{10} is $-OR^9$ where R^9 is selected from hydrogen, C_6 - C_{10} aryl, C_1 - C_8 alkyl, heteroalkyl, $(C_6$ -
15 C_{10} aryl)heteroalkyl, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, $(heteroalkyl)_p(heteroarylene)C_1$ - C_{15} alkylene, $(heteroalkyl)_p(C_6$ - C_{10} arylene) C_1 - C_{15} alkylene, and $(C_1$ - C_{15} alkyl) $_p(C_6$ - C_{10} arylene)heteroalkylene.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^4 and R^5 together form $=NR^{10}$ and R^{10} is
20 $-N(R^9)(R^9)$ where R^9 is selected from hydrogen, C_1 - C_8 alkyl, heteroalkyl, C_6 - C_{10} aryl, $(C_6$ - C_{10} aryl)heteroalkylene, $(heteroalkyl)_pC_6$ - C_{10} arylene, $(C_1$ - C_{15} alkyl) $_pC_6$ - C_{10} arylene, $(heteroalkyl)_p(C_6$ - C_{10} arylene)heteroalkylene, $(C_1$ - C_{15} alkyl) $_p(C_6$ - C_{10} arylene) C_1 - C_{15} alkylene, and $(C_1$ - C_{15} alkyl) $_p(C_6$ - C_{10} arylene) C_1 - C_{15} heteroalkylene.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^4 and R^5 together form $=CR^8R^8$, and one of
25 R^8 is hydrogen while the other R^8 is selected from hydrogen, C_1 - C_8 alkyl and heteroalkyl.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^8 is selected from hydrogen and C_1 - C_8 alkyl, and R^{10} is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from

alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^8 is hydrogen.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^{10} is R^{11} .

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^4 and R^5 together with the carbon to which they are both attached form a spiro carbocyclic or heterocyclic ring.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^6 is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^6 is selected from C_1 - C_{15} alkyl, C_1 - C_{15} heteroalkyl, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, $(C_6$ aryl)(C_6 aryl) C_1 - C_{15} alkylene, $(C_2$ - C_6 heteroaryl) C_1 - C_{15} alkylene, $(C_6$ - C_{10} aryl) C_1 - C_{15} heteroalkylene, $(\text{heteroalkyl})_p(C_6$ - C_{10} arylene) C_1 - C_{15} alkylene, $(\text{heteroalkyl})_p(C_2$ - C_6 heteroarylene) C_1 - C_{15} alkylene, and $(\text{heteroalkyl})_p(C_6$ arylene)(heteroalkylene)(C_6 arylene) C_1 - C_{15} alkylene.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^6 is an inorganic group having 1-8 atoms exclusively selected from boron, sulfur, phosphorous, silicon and hydrogen. Separately, in any of the above embodiments, R^6 excludes inorganic group having 1-8 atoms exclusively selected from boron, sulfur, phosphorous, silicon and hydrogen.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^6 is hydrogen.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment n is 0.


10 In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment n is 1.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^3 is hydrogen; R^4 and R^5 are selected from (a) R^4 is hydrogen and R^5 is hydroxyl or protected hydroxyl and (b) R^4 and R^5 together form carbonyl; R^6 is hydrogen; and n is 0.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^2 is $-OR^9$.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^2 is $-OCH_2CH_2Si(CH_3)_3$.


20 In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^1 is



In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^9 is a C_1 - C_6 hydrocarbyl.

25 In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^9 is selected from n-propyl and $-\text{CH}_2-\text{CH}=\text{CH}_2$.


In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment R^1 is




In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment, R^8 is hydrogen and R^9 is C_1 - C_6 hydrocarbyl.

- 5 In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment, R^9 is $-\text{CH}_2-\text{CH}=\text{CH}_2$.

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment the stereochemistry of the R^1 and $\text{C}(=\text{O})\text{R}^2$ groups being as shown in formula Ia, with R^1 and $\text{C}(=\text{O})\text{R}^2$ in a *cis* arrangement, both over
10 the benzo ring substituted with $-\text{OR}^6$




In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment the stereochemistry of the R^1 and $\text{C}(=\text{O})\text{R}^2$ groups being as shown in formula Ib, with R^1 and $\text{C}(=\text{O})\text{R}^2$ in a *trans* arrangement, with only $\text{C}(=\text{O})\text{R}^2$ over the benzo ring substituted with $-\text{OR}^6$




(Ib).

In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment the stereochemistry of the R^1 and $C(=O)R^2$ groups being as shown in formula Ic, with R^1 and $C(=O)R^2$ in a *trans* arrangement, with only R^1 over the benzo ring substituted with $-OR^6$



(Ic).

- 5 In any of the above embodiments unless otherwise inconsistent with a previous definition, in a further embodiment the stereochemistry of the R^1 and $C(=O)R^2$ groups being as shown in formula Id, with R^1 and $C(=O)R^2$ in a *cis* arrangement, with neither of the R^1 nor $C(=O)R^2$ groups being over the benzo ring substituted with $-OR^6$



(Id).

- 10 In another embodiment, the present invention provides a composition comprising a compound, or a combination of compounds, according to any one of the

above-described embodiments, and a pharmaceutically acceptable carrier, adjuvant or incipient.


5 In another embodiment, the present invention provides a method for inhibiting a TNF- α mediated processes, comprising administering to a patient in need thereof, through a therapeutically or prophylactically acceptable manner, a therapeutically or pharmaceutically effective amount of a composition comprising a compound or a mixture of compounds according to any of the above-described embodiments. In one embodiment, the administering is selected from transdermal, oral, intravenous, intramuscular, vaginal, rectal, pulmonary, subcutaneous, sublingual and transmucosal administration.

10 In another embodiment, the present invention provides a method for treating an inflammation event, comprising administering to a patient in need thereof, through a therapeutically or prophylactically acceptable manner, a therapeutically or pharmaceutically effective amount of a composition comprising a compound or a mixture of compounds according to any of the above-described embodiments. In one embodiment,
15 the administering is selected from transdermal, oral, intravenous, intramuscular, vaginal, rectal, pulmonary, subcutaneous, sublingual and transmucosal administration.

In another embodiment, the present invention provides a library of benzobicyclooctane compounds where said library comprises a plurality of compounds each having a structure of formula (I) as describe above, including inventive embodiments
20 thereof as set forth above, where diversity is present among the R¹, R², R³, R⁴, R⁵, R⁶, and R⁷ groups.

In another embodiment, the present invention provides a process for preparing a combinatorial library of benzobicyclooctane compounds, wherein said library comprises a plurality of compounds of formula (I), including inventive embodiments
25 thereof as set forth above, said process comprising the steps:


(a) providing a compound bound to a solid support according to formula (II)



wherein PG1 and PG2 refer to first and second protecting groups, respectively, where the first protecting group can be removed in the continued presence of the second protecting group, and the second protecting group can be removed in the continued presence of the linker, and (SS) refers to a solid support;

(b) removing the first protecting group but not the second protecting group, to provide a first deprotected product;


(c) reacting the first deprotected product with a plurality of amines of the formula HNR' to provide a plurality of compounds bound to a solid support, each according to formula (IIa)



where R and R' are each independently selected from R^9 ;


(d) removing the second protecting group from (IIa) to provide a second deprotected product;

(e) reacting the second deprotected product with a plurality of amines of the formula $\text{HNR}''\text{R}'''$ to provide a plurality of compounds bound to a solid support, each according to formula (IIb)



where R'' and R''' are each independently selected from R^9 ;

(f) removing the scaffold from the linker to provide a library of compounds according to formula (IIc)



5

These and other embodiments of the present invention are described in further detail below.

DETAILED DESCRIPTION OF THE INVENTION

Before providing a detailed description of the invention, a number of terms as used herein are defined as follows:

10

Definition of terms

As used herein, the following terms have the indicated meanings.

Unless otherwise indicated, the term "a" refers to one or more than one of the indicated items. For example, "a compound" includes one and more than one compound.

15

"Alkyl" is a saturated or unsaturated, straight or branched, hydrocarbon chain. In various embodiments, the alkyl group has 1-18 carbon atoms, *i.e.*, is a C1-C18

group, or is a C1-C12 group, a C1-C6 group, or a C1-C4 group. Independently, in various embodiments, the alkyl group has zero branches (*i.e.*, is a straight chain), one branch, two branches, or more than two branches. Independently, in one embodiment, the alkyl group is saturated. In another embodiment, the alkyl group is unsaturated. In various
5 embodiments, the unsaturated alkyl may have one double bond, two double bonds, more than two double bonds, and/or one triple bond, two triple bonds, or more than two triple bonds. Alkyl chains may be substituted or unsubstituted. In one embodiment, the alkyl chains are unsubstituted. In another embodiment, the alkyl chain is substituted, *e.g.*, with 1 substituent (*i.e.*, the alkyl group is monosubstituted), or 1-2 substituents, or 1-3
10 substituents, or 1-4 substituents, etc.

"Aryl" is an aromatic hydrocarbon ring system. The ring system may be monocyclic or fused polycyclic (*e.g.*, bicyclic, tricyclic, etc.). In various embodiments, the monocyclic aryl ring is C5-C10, or C5-C7, or C5-C6, where these carbon numbers refer to the number of carbon atoms that make up the ring system. A C6 ring system, *i.e.*, a phenyl
15 ring, is a preferred aryl ring. In various embodiments, the polycyclic ring is a bicyclic aryl ring, where preferred bicyclic aryl rings are C8-C12, or C9-C10. A naphthyl ring, which has 10 carbon atoms, is a preferred polycyclic aryl ring. Aryl rings may be substituted or unsubstituted. In one embodiment, the aryl ring is unsubstituted. In another embodiment, the aryl ring is substituted with 1 substituent (*i.e.*, the aryl ring is monosubstituted), or 1-2
20 substituents, or 1-3 substituents, or 1-4 substituents, etc.

"Carbocyclic aliphatic ring," also referred to as carbocycle, is a saturated or unsaturated, monocyclic or polycyclic (*e.g.*, bicyclic, tricyclic, etc.) hydrocarbon ring. Carbocyclic aliphatic rings are not aromatic. A polycyclic hydrocarbon ring may include fused, spiro or bridged ring structures. In various embodiments, the monocyclic
25 carbocyclic aliphatic ring is a C3-C10, or a C4-C7, or a C5-C6 ring system. In various embodiments, the polycyclic carbocyclic aliphatic ring is a C6-C12, or a C9-C10 ring system. In one embodiment, the polycyclic ring is bicyclic. In another embodiment, the polycyclic ring is bicyclic or tricyclic. Carbocyclic aliphatic rings include cyclopropyl, cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl, and cyclooctyl. Carbocycles may be

substituted or unsubstituted. In one embodiment, the carbocycle is unsubstituted. In another embodiment, the carbocycle is substituted with, *e.g.*, 1 substituent (*i.e.*, the alkyl group is monosubstituted), or 1-2 substituents, or 1-3 substituents, or 1-4 substituents, etc.

5 “Haloalkyl” is an alkyl chain substituted with one or more halogens. A preferred haloalkyl is trifluoromethyl.

“Heteroalkyl” is a saturated or unsaturated, straight or branched, chain containing carbon and at least one heteroatom. The heteroalkyl group may, in various embodiments, have one heteroatom, or 1-2 heteroatoms, or 1-3 heteroatoms, or 1-4 heteroatoms. Heteroalkyl chains may contain from 1 to 18 (*i.e.*, 1-18) member atoms
10 (carbon and heteroatoms) in the chain, and in various embodiments contain 1-12, or 1-6, or 1-4 member atoms. Independently, in various embodiments, the heteroalkyl group has zero branches (*i.e.*, is a straight chain), one branch, two branches, or more than two branches. Independently, in one embodiment, the heteroalkyl group is saturated. In another embodiment, the heteroalkyl group is unsaturated. In various embodiments, the
15 unsaturated heteroalkyl may have one double bond, two double bonds, more than two double bonds, and/or one triple bond, two triple bonds, or more than two triple bonds. Heteroalkyl chains may be substituted or unsubstituted. In one embodiment, the heteroalkyl chain is unsubstituted. In another embodiment, the heteroalkyl chain is substituted. A substituted heteroalkyl chain may have 1 substituent (*i.e.*, be
20 monosubstituted), or may have 1-2 substituents, or 1-3 substituents, or 1-4 substituents, etc. Exemplary heteroalkyl groups include esters ($-C(=O)-OR$) and ketones ($-C(=O)-$).

“Heteroaryl” is an aromatic ring system or a semi-aromatic system of rings or a pseudo aromatic ring or rings containing carbon and at least one heteroatom in at least one of the rings. The heteroaryl group may, in various embodiments, have one heteroatom,
25 or 1-2 heteroatoms, or 1-3 heteroatoms, or 1-4 heteroatoms in the ring. The heteroaryl group may further include more than one ring system, which in various embodiments may include one heteroatom or 1-2 heteroatoms, or 1-3 heteroatoms, or 1 heteroatom in each ring system, or 1-4 heteroatoms in each ring system. The heteroaryl group which comprises more than one ring system may, in various embodiments have one or more than

one of the ring systems aromatic. Heteroaryl rings may be monocyclic or polycyclic, where the polycyclic ring may contained fused, spiro or bridged ring junctions. In one embodiment, the heteroaryl is selected from monocyclic and bicyclic. Monocyclic heteroaryl rings may contain from about 5 to about 10 member atoms (carbon and heteroatoms), preferably from 5-7, and most preferably from 5-6 member atoms in the ring. Bicyclic heteroaryl rings may contain from about 8-12 member atoms, or 9-10 member atoms in the ring. The heteroaryl ring may be unsubstituted or substituted. In one embodiment, the heteroaryl ring is unsubstituted. In another embodiment, the heteroaryl ring is substituted. The substituted heteroaryl ring may contain 1 substituent, or 1-2 substituents, or 1-3 substituents, or 1-4 substituents, etc. Exemplary heteroaryl rings include benzofuran, benzothiophene, furan, imidazole, indole, isothiazole, oxazole, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, quinoline, thiazole and thiophene.

"Heteroatom" is a nitrogen, sulfur, oxygen or silicon atom. Groups containing more than one heteroatom may contain different heteroatoms.

"Heterocyclic aliphatic ring," also referred to as "heterocyclyl", is a saturated or unsaturated, monocyclic or polycyclic (e.g., bicyclic, tricyclic, etc.) ring containing carbon and at least one heteroatom. Heterocyclic aliphatic rings are not aromatic *per se* but may be pseudo-aromatic and/or readily be made aromatic through methods known in the art. The heterocyclic aliphatic ring may, in various embodiments, have one heteroatom, or 1-2 heteroatoms, or 1-3 heteroatoms, or 1-4 heteroatoms, etc. In one embodiment, the heterocyclic aliphatic ring is monocyclic, where the monocyclic ring may have 3-10, or 4-7, or 5-6 member atoms. In another embodiment, the heterocyclic aliphatic ring is polycyclic, where in various embodiments, the ring may be bicyclic, or may be tricyclic, or may be either bicyclic or tricyclic. A polycyclic ring system may have one or more fused, spiro or bridged ring systems. The polycyclic heterocyclic aliphatic ring system may have 6-12, or 9-10 member atoms. The heterocyclic ring may be unsubstituted or substituted. In one embodiment, the heterocyclic ring is unsubstituted. In another embodiment, the heterocyclic ring is substituted. The substituted heterocyclic ring may contain 1 substituent, or 1-2 substituents, or 1-3 substituents, or 1-4 substituents, etc.

Exemplary heterocyclic aliphatic rings include piperazyl, morpholinyl, tetrahydrofuranyl, tetrahydropyranyl and piperidyl.

“Inorganic groups having 1-8 atoms exclusively selected from boron, sulfur, phosphorous, silicon and hydrogen” refers to, for example, borates, sulfates, phosphates, silicates, and acids thereof.

“Lower alkyl” is an alkyl chain comprised of 1-6, preferably 1-4 carbon atoms.

“Pharmaceutically acceptable salt” and “salts thereof” means organic or inorganic salts of the pharmaceutically important molecule. A pharmaceutically acceptable salt may involve the inclusion of another molecule such as an acetate ion, a succinate ion or other counterion. The counterion may be any organic or inorganic moiety that stabilizes the charge on the parent compound. Furthermore, a pharmaceutically important organic molecule may have more than one charged atom in its structure. Situations where multiple charged atoms are part of the molecule may have multiple counterions. Hence, the molecule of a pharmaceutically acceptable salt may contain one or more than one charged atoms and may also contain, one or more than one counterion. The desired charge distribution is determined according to methods of drug administration. Examples of pharmaceutically acceptable salts are well known in the art but, without limiting the scope of the present invention, exemplary presentations can be found in the Physician’s Desk Reference, The Merck Index, The Pharmacopoeia and Goodman & Gilman’s The Pharmacological Basis of Therapeutics.

“Substituents” replace a hydrogen atom with a non-hydrogen atom on an alkyl, heteroalkyl, aryl, heteroaryl, carbocycle, and/or heterocyclyl group as defined herein. Where the substituent contains a heteroatom, that heteroatom may be at any acceptable oxidation state for that particular atom, *e.g.*, sulfur as part of a substituent may vary from an oxidation state of -2 to +8, and may be part of a complex or chelate as in a sulfoxide a mercapto-phosphine or metal chelated in a thia-crown ether. Suitable substituents that may be located on one or more of these groups include the following: hydroxy, alkoxy (*i.e.*, alkyl-O-, *e.g.*, methoxy, ethoxy, propoxy, butoxy, pentoxy), aryloxy (*e.g.*, phenoxy, chlorophenoxy, tolyloxy, methoxyphenoxy, benzyloxy, alkyloxycarbonylphenoxy,

acyloxyphenoxy), acyloxy (e.g., propionyloxy, benzoyloxy, acetoxy), carbamoyloxy, carboxy, mercapto, alkylthio, acylthio, arylthio (e.g., phenylthio, chlorophenylthio, alkylphenylthio, alkoxyphenylthio, benzylthio, alkyloxycarbonyl-phenylthio), sulfonamido (5 $-N(R^9)SO_2R^9$ or $-SO_2NR^9R^9$), amino (e.g., amino, mono- and di- C_1-C_3 alkanylamino, methylphenylamino, methylbenzylamino, C_1-C_3 alkanylamido, acylamino, carbamamido, ureido, guanidino, nitro, cyano and imino). Moreover, any substituent may have from 1-5 further substituents attached thereto.

“Amino” means a nitrogen atom substituted with up to 4 groups, for instance, 2 or 3 alkyl groups as defined above, or 1 or 2 alkyl groups and a hydrogen group, 10 or with one or two aryl groups and one or two alkyl groups so that the total number of groups is 2 or 3, or with two or three aryl groups, or with two or more hydrogen atoms or with other the substitution required to complete the nitrogen’s valence requirements. “Amino” further includes amino salts where the nitrogen is hypervalent, having four bonds and may or may not have a charge and a counterion. The counterion, when present, may be 15 an external inorganic and/or organic counterion and/or may be an internal counterion. Inorganic counterions include, for example, anions such as halo anions and other non-metal anions. Examples of organic counterions include, for example, anionic organic moieties such as acetate, citrate and other anionic organic moieties. Thus, amino refers to quaternary ammonium groups, tertiary amines and salts thereof, secondary amines and salts 20 thereof, and primary amines and salts thereof.

As used herein and in the appended claims a “library” means a large number of chemical derivatives used in screening for biological activity or other activity. In general a library will have greater than 20 members, preferably the library will have at least 50 members, more preferably the library will have at least 96 members and most preferably 25 the library will have at least 1000 members.

As used herein and in the appended claims “scaffold” means a common chemical structure found within a library of organic compounds. Similarly, within a combinatorial chemical library the scaffold forms the basis for a diverse series of chemical derivatization, additions and subtractions. Importantly, regardless of the extent of the

chemical derivatization performed on the scaffold, the product is within the scope of the combinatorial library.

“Inflammation event” or “inflammation” or “swelling” are synonymous terms that mean an abnormal enlargement of a portion or tissue of an animal. The abnormal enlargement may be the normal, expected result of another event, such as, for example, sepsis, fever, trauma, shock, or injury. Non-limiting examples of some of these events include sepsis due to renal or liver failure, fever secondary to systemic infection, localized fever secondary to local infection, blunt force trauma or emotional trauma having physical manifestations, shock secondary to trauma and/or other events causing a pooling of body fluids and an injury causing release of cellular fluids initiating the inflammation cascade.

As used herein, “commercially available chemicals” and the chemicals used in the Examples set forth herein may be obtained from standard commercial sources including Acros Organics (Pittsburgh PA), Aldrich Chemical (Milwaukee WI, including Sigma Chemical and Fluka), Apin Chemicals Ltd. (Milton Park UK), Avocado Research (Lancashire U.K.), BDH Inc. (Toronto, Canada), Bionet (Cornwall, U.K.), Chemservice Inc. (West Chester PA), Crescent Chemical Co. (Hauppauge NY), Eastman Organic Chemicals, Eastman Kodak Company (Rochester NY), Fisher Scientific Co. (Pittsburgh PA), Fisons Chemicals (Leicestershire UK), Frontier Scientific (Logan UT), ICN Biomedicals, Inc. (Costa Mesa CA), Key Organics (Cornwall U.K.), Lancaster Synthesis (Windham NH), Maybridge Chemical Co. Ltd. (Cornwall U.K.), Parish Chemical Co. (Orem UT), Pfaltz & Bauer, Inc. (Waterbury CN), Polyorganix (Houston TX), Pierce Chemical Co. (Rockford IL), Riedel de Haen AG (Hannover, Germany), Spectrum Quality Product, Inc. (New Brunswick, NJ), TCI America (Portland OR), Trans World Chemicals, Inc. (Rockville MD), and Wako Chemicals USA, Inc. (Richmond VA).

As used herein, “compounds described in the chemical literature” may be identified through various reference books and databases. Suitable reference books and treatise that detail the synthesis of reactants useful in the preparation of compounds of the present invention, or provide references to articles that describe the preparation, include for example, "Synthetic Organic Chemistry", John Wiley & Sons, Inc., New York; S. R.


Sandler et al., "Organic Functional Group Preparations," 2nd Ed., Academic Press, New York, 1983; H. O. House, "Modern Synthetic Reactions", 2nd Ed., W. A. Benjamin, Inc. Menlo Park, Calif. 1972; T. L. Gilchrist, "Heterocyclic Chemistry", 2nd Ed., John Wiley & Sons, New York, 1992; J. March, "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th Ed., Wiley-Interscience, New York, 1992. Specific and analogous reactants may also be identified through the indices of known chemicals prepared by the Chemical Abstract Service of the American Chemical Society, which are available in most public and university libraries, as well as through on-line databases (the American Chemical Society, Washington, D.C., www.acs.org may be contacted for more details). Chemicals that are known but not commercially available in catalogs may be prepared by custom chemical synthesis houses, where many of the standard chemical supply houses (e.g., those listed above) provide custom synthesis services.

As used herein "suitable conditions" for carrying out a synthetic step are explicitly provided herein or may be discerned by reference to publications directed to methods used in synthetic organic chemistry. The reference books and treatise set forth above that detail the synthesis of reactants useful in the preparation of compounds of the present invention, will also provide suitable conditions for carrying out a synthetic step according to the present invention.

All other acronyms and abbreviations have the corresponding meaning as published in journals relative to the art of organic chemistry.

A. Compounds

In one aspect, the present invention provides benzobicyclooctane compounds of formula (I)



(I)

and optical isomers, diastereomers, enantiomers and pharmaceutically acceptable salts thereof in isolation (*i.e.*, isolated from one another) or in mixture (*i.e.*, two or more compounds in admixture with one another), where, independently at each location:

R^1 is selected from the following six formulae:

R^2 is $-OR^9$ or $-NR^9R^9$;

10 R^3 is selected from hydrogen, halogen, hydroxyl or protected hydroxyl, amino or protected amino, and C_1 - C_8 alkyl or C_1 - C_8 haloalkyl;

R^4 and R^5 are independently selected from R^9 , $-OR^9$, $-NR^9R^9$ and $-N=N-R^9$, or R^4 and R^5 may together form a group selected from $=O$, $=CR^8R^8$ and $=NR^{10}$, or R^4 and R^5 may together with the carbon to which they are both attached form a spiro carbocyclic
15 or heterocyclic ring;

R^6 is selected from hydrogen, inorganic groups having 1-8 atoms exclusively selected from boron, sulfur, phosphorous, silicon and hydrogen, and organic

groups having 1-20 carbons and optionally containing 1-4 heteroatoms selected from nitrogen, oxygen and silicon;

R^7 is selected from halogen, hydroxyl or protected hydroxyl, amino or protected amino, and C_1 - C_8 alkyl or C_1 - C_8 haloalkyl;

5 R^8 is selected from hydrogen, alkyl (preferably C_1 - C_8 alkyl), aryl and heteroalkyl;


R^9 is selected from hydrogen and organic groups having 1-30 carbons and optionally containing 1-4 heteroatoms selected from nitrogen, oxygen and silicon, with the proviso that two R^9 groups both joined to a common atom may be joined together so as to
10 form a ring with the common atom;

R^{10} is selected from $-R^9$, $-OR^9$, $-NR^9R^9$, $-NH-C(O)R^9$; $-NH-C(O)OR^9$ and $-NH-C(S)NHR^9$; and

n is 0, 1, 2 or 3.

In one embodiment, when R^6 is hydrogen and R^4 and R^5 together form $=O$
15 and R^1 is CO_2R^9 , then R^2 is not OCH_3 . In one embodiment, R^4 and R^5 are both hydrogen, while in another embodiment R^4 is not hydrogen when R^5 is hydrogen.

In formula (1), the two wavy lines (one connected to R^1 , the other connected to $C(=O)R^2$) indicate that the invention provides any possible stereochemistry for the R^1 and $C(=O)R^2$ groups. In other words, the present invention provides benzobicyclooctanes having
20 each of the four relative stereochemistries shown below as formulae (Ia), (Ib), (Ic) and (Id).



In individual aspects, the present invention provides compounds of formulae Ia through Io, where each of Ia through Io is made up of one or more of the compounds of formula Ia, Ib, Ic and Id. An "x" in a box to the right of the designation Ia through Io indicates which of Ia, Ib, Ic and Id is contained within the designated formula. Thus, for instance, the compounds of formula Ij contain the compounds within formulae Ic and Id (as an "x" is present in the columns designated Ic and Id), but do not include the compounds of formula Ia or Ib (as no "x" appears in the columns designated Ia and Ib) in the row designated formula Ij.

Table A

Code No.	Formula			
	Ia	Ib	Ic	Id
Ia	x			
Ib		x		
Ic			x	
Id				x
Ie	x	x		
If	x		x	
Ig	x			x
Ih		x	x	







Code No.	Formula			
	Ia	Ib	Ic	Id
Ii		x		x
Ij			x	x
Ik	x	x	x	
Il	x	x		x
Im	x		x	x
In		x	x	x
Io	x	x	x	x

Thus, as shown in Table A, in one aspect the present invention provides compounds of formula Ia, while in a separate aspect the present invention provides compounds of formula Ib; while in a still separate aspect the present invention provides compounds of formula Ic. In another aspect, the present invention provides compounds of formula Id, while in another aspect the present invention provides compounds of formula Ie (containing the set of compounds within formulae Ia and Ib), and in another aspect the present invention provides compounds of formula If (containing the set of compounds within formulae Ia and Ic). In still another aspect the present invention provides compounds of formula Ig, and in another aspect provides compounds of formula Ih, while in another aspect the invention provides compounds of formula Ii, and in yet another aspect the present invention provides compounds of formula Ij. In a separate aspect, the present invention provides compounds of formula Ik, while in another aspect the present invention provides compound of formula Il, and in still another aspect the invention provides compounds of formula Im. In addition, the present invention provides compounds of formula In, while in another aspect the present invention provides compounds of formula

10. Thus, using a convenient shorthand, it may be said that in various aspects the present invention provides benzybicyclooctane compounds of formulae: (Ia); (Ib); (Ic); (Id); (Ie); (If); (Ig); (Ih); (Ii); (Ij); (Ik); (Il); (Im); (In); (Io). In each of the above-listed aspects, the compounds include optical isomers, diastereomers, enantiomers and pharmaceutically acceptable salts thereof in isolation or mixture, where, independently at each location, the substituents R^1 , R^2 etc. are as defined herein.

In the compounds of the present invention, R^1 is selected from the following six formulae, identified as R1a, R1b, R1c, R1d, R1e and R1f as defined below in Table B.

Table B

 <p>R1a</p>	 <p>R1b</p>	 <p>R1c</p>
 <p>R1d</p>	 <p>R1e</p>	 <p>R1f</p>

Thus, in compounds of the invention, R^1 may be one or more of R1a, R1b, R1c, R1d, R1e and R1f. Table C defines groups R1A through R1BJ, where each of R1A through R1BJ is composed of one or more of R1a, R1b, R1c, R1d, R1e and R1f. For example, as shown in Table C, R1A is defined as formula R1a but not any of R1b through R1f. As another example, R1G is defined as the sum of R1a and R1b, but does not include R1c through R1f. As a final example, R1BI is the sum of R1b through R1f, and excludes only R1a.

Table C

	R1					
R1	a	b	c	d	e	f
A	X					
B		X				
C			X			
D				X		
E					X	
F						X
G	X	X				
H	X		X			
I	X			X		
J	X				X	
K	X					X
L		X	X			
M		X		X		
N		X			X	
O		X				X
P			X	X		
Q			X		X	
R			X			X
S				X	X	
T				X		X
U					X	X
V	X	X	X			
W	X	X		X		
X	X	X			X	

	R1					
R1	a	b	c	d	e	f
Y	X	X				X
Z	X		X	X		
AA	X		X		X	
AB	X		X			X
AC	X			X	X	
AD	X			X		X
AE	X				X	X
AF		X	X	X		
AG		X	X		X	
AH		X	X			X
AI		X		X	X	
AJ		X		X		X
AK		X			X	X
AL			X	X	X	
AM			X	X		X
AN			X		X	X
AO				X	X	X
AP	X	X	X	X		
AQ	X	X	X		X	
AR	X	X	X			X
AS	X	X		X	X	
AT	X	X		X		X
AU	X	X			X	X
AV	X		X	X	X	
AW	X		X	X		X
AX	X		X		X	X

	R1					
R1	a	b	c	d	e	f
AY	X			X	X	X
AZ		X	X	X	X	
BA		X	X	X		X
BB		X		X	X	X
BC			X	X	X	X
BD	X	X	X	X	X	
BE	X	X	X	X		X
BF	X	X	X		X	X
BG	X	X		X	X	X
BH	X		X	X	X	X
BI		X	X	X	X	X
BJ	X	X	X	X	X	X

Thus, in one aspect, the present invention provides compounds of formula (I) where R^1 is R1A. In another aspect, the invention provides compounds of formula (I) where R^1 is R1B. In another aspect, the invention provides compounds of formula (I) where R^1 is R1C. In another aspect, the invention provides compounds of formula (I) where R^1 is R1D. In another aspect, the invention provides compounds of formula (I) where R^1 is R1E. In another aspect, the invention provides compounds of formula (I) where R^1 is R1F. In other words, stated in a convenient shorthand nomenclature, in various aspects the present invention provides "Compounds of formula (I) where: R^1 is R1A; R^1 is R1B; R^1 is R1C; R^1 is R1D; R^1 is R1E; R^1 is R1F."

Using this same shorthand nomenclature, in various aspects the present invention provides compounds of formula (I) where: R^1 is R1G; R^1 is R1H; R^1 is R1I; R^1 is R1J; R^1 is R1K; R^1 is R1L; R^1 is R1M; R^1 is R1N; R^1 is R1O; R^1 is R1P; R^1 is R1Q; R^1 is R1R; R^1 is R1S; R^1 is R1T; R^1 is R1U; R^1 is R1V; R^1 is R1W; R^1 is R1X; R^1 is R1Y; R^1

is R1Z; R¹ is R1AA; R¹ is R1AB; R¹ is R1AC; R¹ is R1AD; R¹ is R1AE; R¹ is R1AF; R¹ is R1AG; R¹ is R1AH; R¹ is R1AI; R¹ is R1AJ; R¹ is R1AK; R¹ is R1AL; R¹ is R1AM; R¹ is R1AN; R¹ is R1AO; R¹ is R1AP; R¹ is R1AQ; R¹ is R1AR; R¹ is R1AS; R¹ is R1AT; R¹ is R1AU; R¹ is R1AV; R¹ is R1AW; R¹ is R1AX; R¹ is R1AY; R¹ is R1AZ; R¹ is R1BA; R¹ is R1BB; R¹ is R1BC; R¹ is R1BD; R¹ is R1BE; R¹ is R1BF; R¹ is R1BG; R¹ is R1BH; R¹ is R1BI; R¹ is R1BJ.

In separate aspects, the present invention provides compounds of formulae (Ia)-(Io) as defined in Table B wherein R¹ is selected from R1A through R1BJ as defined in Table C. Each of these aspects is given a unique identifier, X1 through X937 in Table D, where each of X1 through X937 is a separate and unique aspect of the present invention. In each of X1 through X937, R² is -OR⁹ or NR⁹R⁹. The present invention also provides aspects Y1 through Y937 as defined in Table E, which are analogous to aspects X1 through X937 in terms of formula (Ia)-(Io) and R¹, however in aspects Y1 through Y937 R² is limited to -OR⁹. The present invention also provides aspects Z1 through Z937 as defined in Table F, which are analogous to aspects X1 through X937 in terms of formula (Ia)-(Io) and R¹, however in aspects Z1 through Z937 R² is limited to -NR⁹R⁹.

Table D

R1	Ia	Ib	Ic	Id	Ie	If	Ig	Ih	Ii	Ij	Ik	Il	Im	In	Io
A	X1	X63	X125	X187	X249	X311	X373	X435	X497	X559	X621	X683	X745	X807	X869
B	X2	X64	X126	X188	X250	X312	X374	X436	X498	X560	X622	X684	X746	X808	X870
C	X3	X65	X127	X189	X251	X313	X375	X437	X499	X561	X623	X685	X747	X809	X871
D	X4	X66	X128	X190	X252	X314	X376	X438	X500	X562	X624	X686	X748	X810	X872
E	X5	X67	X129	X191	X253	X315	X377	X439	X501	X563	X625	X687	X749	X811	X873
F	X6	X68	X130	X192	X254	X316	X378	X440	X502	X564	X626	X688	X750	X812	X874
G	X7	X69	X131	X193	X255	X317	X379	X441	X503	X565	X627	X689	X751	X813	X875
H	X8	X70	X132	X194	X256	X318	X380	X442	X504	X566	X628	X690	X752	X814	X876
I	X9	X71	X133	X195	X257	X319	X381	X443	X505	X567	X629	X691	X753	X815	X877
J	X10	X72	X134	X196	X258	X320	X382	X444	X506	X568	X630	X692	X754	X816	X878
K	X11	X73	X135	X197	X259	X321	X383	X445	X507	X569	X631	X693	X755	X817	X879
L	X12	X74	X136	X198	X260	X322	X384	X446	X508	X570	X632	X694	X756	X818	X880
M	X13	X75	X137	X199	X261	X323	X385	X447	X509	X571	X633	X695	X757	X819	X881
N	X14	X76	X138	X200	X262	X324	X386	X448	X510	X572	X634	X696	X758	X820	X882
O	X15	X77	X139	X201	X263	X325	X387	X449	X511	X573	X635	X697	X759	X821	X883
P	X16	X78	X140	X202	X264	X326	X388	X450	X512	X574	X636	X698	X760	X822	X884
Q	X17	X79	X141	X203	X265	X327	X389	X451	X513	X575	X637	X699	X761	X823	X885
R	X18	X80	X142	X204	X266	X328	X390	X452	X514	X576	X638	X700	X762	X824	X886
S	X19	X81	X143	X205	X267	X329	X391	X453	X515	X577	X639	X701	X763	X825	X887
T	X20	X82	X144	X206	X268	X330	X392	X454	X516	X578	X640	X702	X764	X826	X888
U	X21	X83	X145	X207	X269	X331	X393	X455	X517	X579	X641	X703	X765	X827	X889
V	X22	X84	X146	X208	X270	X332	X394	X456	X518	X580	X642	X704	X766	X828	X890
W	X23	X85	X147	X209	X271	X333	X395	X457	X519	X581	X643	X705	X767	X829	X891
X	X24	X86	X148	X210	X272	X334	X396	X458	X520	X582	X644	X706	X768	X830	X892
Y	X25	X87	X149	X211	X273	X335	X397	X459	X521	X583	X645	X707	X769	X831	X893
Z	X26	X88	X150	X212	X274	X336	X398	X460	X522	X584	X646	X708	X770	X832	X894

Table D (cont)

AA	X27	X89	X151	X213	X275	X337	X399	X461	X523	X585	X647	X709	X771	X833	X895
AB	X28	X90	X152	X214	X276	X338	X400	X462	X524	X586	X648	X710	X772	X834	X896
AC	X29	X91	X153	X215	X277	X339	X401	X463	X525	X587	X649	X711	X773	X835	X897
AD	X30	X92	X154	X216	X278	X340	X402	X464	X526	X588	X650	X712	X774	X836	X898
AE	X31	X93	X155	X217	X279	X341	X403	X465	X527	X589	X651	X713	X775	X837	X899
AF	X32	X94	X156	X218	X280	X342	X404	X466	X528	X590	X652	X714	X776	X838	X900
AG	X33	X95	X157	X219	X281	X343	X405	X467	X529	X591	X653	X715	X777	X839	X901
AH	X34	X96	X158	X220	X282	X344	X406	X468	X530	X592	X654	X716	X778	X840	X902
AI	X35	X97	X159	X221	X283	X345	X407	X469	X531	X593	X655	X717	X779	X841	X903
AJ	X36	X98	X160	X222	X284	X346	X408	X470	X532	X594	X656	X718	X780	X842	X904
AK	X37	X99	X161	X223	X285	X347	X409	X471	X533	X595	X657	X719	X781	X843	X905
AL	X38	X100	X162	X224	X286	X348	X410	X472	X534	X596	X658	X720	X782	X844	X906
AM	X39	X101	X163	X225	X287	X349	X411	X473	X535	X597	X659	X721	X783	X845	X907
AN	X40	X102	X164	X226	X288	X350	X412	X474	X536	X598	X660	X722	X784	X846	X908
AO	X41	X103	X165	X227	X289	X351	X413	X475	X537	X599	X661	X723	X785	X847	X909
AP	X42	X104	X166	X228	X290	X352	X414	X476	X538	X600	X662	X724	X786	X848	X910
AW	X43	X105	X167	X229	X291	X353	X415	X477	X539	X601	X663	X725	X787	X849	X911
AR	X44	X106	X168	X230	X292	X354	X416	X478	X540	X602	X664	X726	X788	X850	X912
AS	X45	X107	X169	X231	X293	X355	X417	X479	X541	X603	X665	X727	X789	X851	X913
AT	X46	X108	X170	X232	X294	X356	X418	X480	X542	X604	X666	X728	X790	X852	X914
AU	X47	X109	X171	X233	X295	X357	X419	X481	X543	X605	X667	X729	X791	X853	X915
AV	X48	X110	X172	X234	X296	X358	X420	X482	X544	X606	X668	X730	X792	X854	X916
AW	X49	X111	X173	X235	X297	X359	X421	X483	X545	X607	X669	X731	X793	X855	X917
AX	X50	X112	X174	X236	X298	X360	X422	X484	X546	X608	X670	X732	X794	X856	X918
AY	X51	X113	X175	X237	X299	X361	X423	X485	X547	X609	X671	X733	X795	X857	X919
AZ	X52	X114	X176	X238	X300	X362	X424	X486	X548	X610	X672	X734	X796	X858	X920

Table D (cont)

BA	X53	X115	X177	X239	X301	X363	X425	X487	X549	X611	X673	X735	X797	X859	X921
BB	X54	X116	X178	X240	X302	X364	X426	X488	X550	X612	X674	X736	X798	X860	X922
BC	X55	X117	X179	X241	X303	X365	X427	X489	X551	X613	X675	X737	X799	X861	X923
BD	X56	X118	X180	X242	X304	X366	X428	X490	X552	X614	X676	X738	X800	X862	X924
BE	X57	X119	X181	X243	X305	X367	X429	X491	X553	X615	X677	X739	X801	X863	X925
BF	X58	X120	X182	X244	X306	X368	X430	X492	X554	X616	X678	X740	X802	X864	X926
BG	X59	X121	X183	X245	X307	X369	X431	X493	X555	X617	X679	X741	X803	X865	X927
BH	X60	X122	X184	X246	X308	X370	X432	X494	X556	X618	X680	X742	X804	X866	X928
BI	X61	X123	X185	X247	X309	X371	X433	X495	X557	X619	X681	X743	X805	X867	X929
BJ	X62	X124	X186	X248	X310	X372	X434	X496	X558	X620	X682	X744	X806	X868	X930

Table E

R1	Ia	Ib	Ic	Id	Ie	If	Ig	Ih	Ii	Ij	Ik	Il	Im	In	Io
A	Y1	Y63	Y125	Y187	Y249	Y311	Y373	Y435	Y497	Y559	Y621	Y683	Y745	Y807	Y869
B	Y2	Y64	Y126	Y188	Y250	Y312	Y374	Y436	Y498	Y560	Y622	Y684	Y746	Y808	Y870
C	Y3	Y65	Y127	Y189	Y251	Y313	Y375	Y437	Y499	Y561	Y623	Y685	Y747	Y809	Y871
D	Y4	Y66	Y128	Y190	Y252	Y314	Y376	Y438	Y500	Y562	Y624	Y686	Y748	Y810	Y872
E	Y5	Y67	Y129	Y191	Y253	Y315	Y377	Y439	Y501	Y563	Y625	Y687	Y749	Y811	Y873
F	Y6	Y68	Y130	Y192	Y254	Y316	Y378	Y440	Y502	Y564	Y626	Y688	Y750	Y812	Y874
G	Y7	Y69	Y131	Y193	Y255	Y317	Y379	Y441	Y503	Y565	Y627	Y689	Y751	Y813	Y875
H	Y8	Y70	Y132	Y194	Y256	Y318	Y380	Y442	Y504	Y566	Y628	Y690	Y752	Y814	Y876
I	Y9	Y71	Y133	Y195	Y257	Y319	Y381	Y443	Y505	Y567	Y629	Y691	Y753	Y815	Y877
J	Y10	Y72	Y134	Y196	Y258	Y320	Y382	Y444	Y506	Y568	Y630	Y692	Y754	Y816	Y878
K	Y11	Y73	Y135	Y197	Y259	Y321	Y383	Y445	Y507	Y569	Y631	Y693	Y755	Y817	Y879

Table E (cont)

L	Y12	Y74	Y136	Y198	Y260	Y322	Y384	Y446	Y508	Y570	Y632	Y694	Y756	Y818	Y880
M	Y13	Y75	Y137	Y199	Y261	Y323	Y385	Y447	Y509	Y571	Y633	Y695	Y757	Y819	Y881
N	Y14	Y76	Y138	Y200	Y262	Y324	Y386	Y448	Y510	Y572	Y634	Y696	Y758	Y820	Y882
O	Y15	Y77	Y139	Y201	Y263	Y325	Y387	Y449	Y511	Y573	Y635	Y697	Y759	Y821	Y883
P	Y16	Y78	Y140	Y202	Y264	Y326	Y388	Y450	Y512	Y574	Y636	Y698	Y760	Y822	Y884
Q	Y17	Y79	Y141	Y203	Y265	Y327	Y389	Y451	Y513	Y575	Y637	Y699	Y761	Y823	Y885
R	Y18	Y80	Y142	Y204	Y266	Y328	Y390	Y452	Y514	Y576	Y638	Y700	Y762	Y824	Y886
S	Y19	Y81	Y143	Y205	Y267	Y329	Y391	Y453	Y515	Y577	Y639	Y701	Y763	Y825	Y887
T	Y20	Y82	Y144	Y206	Y268	Y330	Y392	Y454	Y516	Y578	Y640	Y702	Y764	Y826	Y888
U	Y21	Y83	Y145	Y207	Y269	Y331	Y393	Y455	Y517	Y579	Y641	Y703	Y765	Y827	Y889
V	Y22	Y84	Y146	Y208	Y270	Y332	Y394	Y456	Y518	Y580	Y642	Y704	Y766	Y828	Y890
W	Y23	Y85	Y147	Y209	Y271	Y333	Y395	Y457	Y519	Y581	Y643	Y705	Y767	Y829	Y891
X	Y24	Y86	Y148	Y210	Y272	Y334	Y396	Y458	Y520	Y582	Y644	Y706	Y768	Y830	Y892
Y	Y25	Y87	Y149	Y211	Y273	Y335	Y397	Y459	Y521	Y583	Y645	Y707	Y769	Y831	Y893
Z	Y26	Y88	Y150	Y212	Y274	Y336	Y398	Y460	Y522	Y584	Y646	Y708	Y770	Y832	Y894
AA	Y27	Y89	Y151	Y213	Y275	Y337	Y399	Y461	Y523	Y585	Y647	Y709	Y771	Y833	Y895
AB	Y28	Y90	Y152	Y214	Y276	Y338	Y400	Y462	Y524	Y586	Y648	Y710	Y772	Y834	Y896
AC	Y29	Y91	Y153	Y215	Y277	Y339	Y401	Y463	Y525	Y587	Y649	Y711	Y773	Y835	Y897
AD	Y30	Y92	Y154	Y216	Y278	Y340	Y402	Y464	Y526	Y588	Y650	Y712	Y774	Y836	Y898
AE	Y31	Y93	Y155	Y217	Y279	Y341	Y403	Y465	Y527	Y589	Y651	Y713	Y775	Y837	Y899
AF	Y32	Y94	Y156	Y218	Y280	Y342	Y404	Y466	Y528	Y590	Y652	Y714	Y776	Y838	Y900
AG	Y33	Y95	Y157	Y219	Y281	Y343	Y405	Y467	Y529	Y591	Y653	Y715	Y777	Y839	Y901
AH	Y34	Y96	Y158	Y220	Y282	Y344	Y406	Y468	Y530	Y592	Y654	Y716	Y778	Y840	Y902
AI	Y35	Y97	Y159	Y221	Y283	Y345	Y407	Y469	Y531	Y593	Y655	Y717	Y779	Y841	Y903
AJ	Y36	Y98	Y160	Y222	Y284	Y346	Y408	Y470	Y532	Y594	Y656	Y718	Y780	Y842	Y904
AK	Y37	Y99	Y161	Y223	Y285	Y347	Y409	Y471	Y533	Y595	Y657	Y719	Y781	Y843	Y905

Table E (cont)

AL	Y38	Y100	Y162	Y224	Y286	Y348	Y410	Y472	Y534	Y596	Y658	Y720	Y782	Y844	Y906
AM	Y39	Y101	Y163	Y225	Y287	Y349	Y411	Y473	Y535	Y597	Y659	Y721	Y783	Y845	Y907
AN	Y40	Y102	Y164	Y226	Y288	Y350	Y412	Y474	Y536	Y598	Y660	Y722	Y784	Y846	Y908
AO	Y41	Y103	Y165	Y227	Y289	Y351	Y413	Y475	Y537	Y599	Y661	Y723	Y785	Y847	Y909
AP	Y42	Y104	Y166	Y228	Y290	Y352	Y414	Y476	Y538	Y600	Y662	Y724	Y786	Y848	Y910
AW	Y43	Y105	Y167	Y229	Y291	Y353	Y415	Y477	Y539	Y601	Y663	Y725	Y787	Y849	Y911
AR	Y44	Y106	Y168	Y230	Y292	Y354	Y416	Y478	Y540	Y602	Y664	Y726	Y788	Y850	Y912
AS	Y45	Y107	Y169	Y231	Y293	Y355	Y417	Y479	Y541	Y603	Y665	Y727	Y789	Y851	Y913
AT	Y46	Y108	Y170	Y232	Y294	Y356	Y418	Y480	Y542	Y604	Y666	Y728	Y790	Y852	Y914
AU	Y47	Y109	Y171	Y233	Y295	Y357	Y419	Y481	Y543	Y605	Y667	Y729	Y791	Y853	Y915
AV	Y48	Y110	Y172	Y234	Y296	Y358	Y420	Y482	Y544	Y606	Y668	Y730	Y792	Y854	Y916
AW	Y49	Y111	Y173	Y235	Y297	Y359	Y421	Y483	Y545	Y607	Y669	Y731	Y793	Y855	Y917
AX	Y50	Y112	Y174	Y236	Y298	Y360	Y422	Y484	Y546	Y608	Y670	Y732	Y794	Y856	Y918
AY	Y51	Y113	Y175	Y237	Y299	Y361	Y423	Y485	Y547	Y609	Y671	Y733	Y795	Y857	Y919
AZ	Y52	Y114	Y176	Y238	Y300	Y362	Y424	Y486	Y548	Y610	Y672	Y734	Y796	Y858	Y920
BA	Y53	Y115	Y177	Y239	Y301	Y363	Y425	Y487	Y549	Y611	Y673	Y735	Y797	Y859	Y921
BB	Y54	Y116	Y178	Y240	Y302	Y364	Y426	Y488	Y550	Y612	Y674	Y736	Y798	Y860	Y922
BC	Y55	Y117	Y179	Y241	Y303	Y365	Y427	Y489	Y551	Y613	Y675	Y737	Y799	Y861	Y923
BD	Y56	Y118	Y180	Y242	Y304	Y366	Y428	Y490	Y552	Y614	Y676	Y738	Y800	Y862	Y924
BE	Y57	Y119	Y181	Y243	Y305	Y367	Y429	Y491	Y553	Y615	Y677	Y739	Y801	Y863	Y925
BF	Y58	Y120	Y182	Y244	Y306	Y368	Y430	Y492	Y554	Y616	Y678	Y740	Y802	Y864	Y926
BG	Y59	Y121	Y183	Y245	Y307	Y369	Y431	Y493	Y555	Y617	Y679	Y741	Y803	Y865	Y927
BH	Y60	Y122	Y184	Y246	Y308	Y370	Y432	Y494	Y556	Y618	Y680	Y742	Y804	Y866	Y928
BI	Y61	Y123	Y185	Y247	Y309	Y371	Y433	Y495	Y557	Y619	Y681	Y743	Y805	Y867	Y929
BJ	Y62	Y124	Y186	Y248	Y310	Y372	Y434	Y496	Y558	Y620	Y682	Y744	Y806	Y868	Y930

Table F

R1	Ia	Ib	Ic	Id	Ie	If	Ig	Ih	Ii	Ij	Ik	Il	Im	In	Io
A	Z1	Z63	Z125	Z187	Z249	Z311	Z373	Z435	Z497	Z559	Z621	Z683	Z745	Z807	Z869
B	Z2	Z64	Z126	Z188	Z250	Z312	Z374	Z436	Z498	Z560	Z622	Z684	Z746	Z808	Z870
C	Z3	Z65	Z127	Z189	Z251	Z313	Z375	Z437	Z499	Z561	Z623	Z685	Z747	Z809	Z871
D	Z4	Z66	Z128	Z190	Z252	Z314	Z376	Z438	Z500	Z562	Z624	Z686	Z748	Z810	Z872
E	Z5	Z67	Z129	Z191	Z253	Z315	Z377	Z439	Z501	Z563	Z625	Z687	Z749	Z811	Z873
F	Z6	Z68	Z130	Z192	Z254	Z316	Z378	Z440	Z502	Z564	Z626	Z688	Z750	Z812	Z874
G	Z7	Z69	Z131	Z193	Z255	Z317	Z379	Z441	Z503	Z565	Z627	Z689	Z751	Z813	Z875
H	Z8	Z70	Z132	Z194	Z256	Z318	Z380	Z442	Z504	Z566	Z628	Z690	Z752	Z814	Z876
I	Z9	Z71	Z133	Z195	Z257	Z319	Z381	Z443	Z505	Z567	Z629	Z691	Z753	Z815	Z877
J	Z10	Z72	Z134	Z196	Z258	Z320	Z382	Z444	Z506	Z568	Z630	Z692	Z754	Z816	Z878
K	Z11	Z73	Z135	Z197	Z259	Z321	Z383	Z445	Z507	Z569	Z631	Z693	Z755	Z817	Z879
L	Z12	Z74	Z136	Z198	Z260	Z322	Z384	Z446	Z508	Z570	Z632	Z694	Z756	Z818	Z880
M	Z13	Z75	Z137	Z199	Z261	Z323	Z385	Z447	Z509	Z571	Z633	Z695	Z757	Z819	Z881
N	Z14	Z76	Z138	Z200	Z262	Z324	Z386	Z448	Z510	Z572	Z634	Z696	Z758	Z820	Z882
O	Z15	Z77	Z139	Z201	Z263	Z325	Z387	Z449	Z511	Z573	Z635	Z697	Z759	Z821	Z883
P	Z16	Z78	Z140	Z202	Z264	Z326	Z388	Z450	Z512	Z574	Z636	Z698	Z760	Z822	Z884
Q	Z17	Z79	Z141	Z203	Z265	Z327	Z389	Z451	Z513	Z575	Z637	Z699	Z761	Z823	Z885
R	Z18	Z80	Z142	Z204	Z266	Z328	Z390	Z452	Z514	Z576	Z638	Z700	Z762	Z824	Z886
S	Z19	Z81	Z143	Z205	Z267	Z329	Z391	Z453	Z515	Z577	Z639	Z701	Z763	Z825	Z887
T	Z20	Z82	Z144	Z206	Z268	Z330	Z392	Z454	Z516	Z578	Z640	Z702	Z764	Z826	Z888
U	Z21	Z83	Z145	Z207	Z269	Z331	Z393	Z455	Z517	Z579	Z641	Z703	Z765	Z827	Z889
V	Z22	Z84	Z146	Z208	Z270	Z332	Z394	Z456	Z518	Z580	Z642	Z704	Z766	Z828	Z890
W	Z23	Z85	Z147	Z209	Z271	Z333	Z395	Z457	Z519	Z581	Z643	Z705	Z767	Z829	Z891
X	Z24	Z86	Z148	Z210	Z272	Z334	Z396	Z458	Z520	Z582	Z644	Z706	Z768	Z830	Z892
Y	Z25	Z87	Z149	Z211	Z273	Z335	Z397	Z459	Z521	Z583	Z645	Z707	Z769	Z831	Z893
Z	Z26	Z88	Z150	Z212	Z274	Z336	Z398	Z460	Z522	Z584	Z646	Z708	Z770	Z832	Z894

Table F (cont)


AA	Z27	Z89	Z151	Z213	Z275	Z337	Z399	Z461	Z523	Z585	Z647	Z709	Z771	Z833	Z895
AB	Z28	Z90	Z152	Z214	Z276	Z338	Z400	Z462	Z524	Z586	Z648	Z710	Z772	Z834	Z896
AC	Z29	Z91	Z153	Z215	Z277	Z339	Z401	Z463	Z525	Z587	Z649	Z711	Z773	Z835	Z897
AD	Z30	Z92	Z154	Z216	Z278	Z340	Z402	Z464	Z526	Z588	Z650	Z712	Z774	Z836	Z898
AE	Z31	Z93	Z155	Z217	Z279	Z341	Z403	Z465	Z527	Z589	Z651	Z713	Z775	Z837	Z899
AF	Z32	Z94	Z156	Z218	Z280	Z342	Z404	Z466	Z528	Z590	Z652	Z714	Z776	Z838	Z900
AG	Z33	Z95	Z157	Z219	Z281	Z343	Z405	Z467	Z529	Z591	Z653	Z715	Z777	Z839	Z901
AH	Z34	Z96	Z158	Z220	Z282	Z344	Z406	Z468	Z530	Z592	Z654	Z716	Z778	Z840	Z902
AI	Z35	Z97	Z159	Z221	Z283	Z345	Z407	Z469	Z531	Z593	Z655	Z717	Z779	Z841	Z903
AJ	Z36	Z98	Z160	Z222	Z284	Z346	Z408	Z470	Z532	Z594	Z656	Z718	Z780	Z842	Z904
AK	Z37	Z99	Z161	Z223	Z285	Z347	Z409	Z471	Z533	Z595	Z657	Z719	Z781	Z843	Z905
AL	Z38	Z100	Z162	Z224	Z286	Z348	Z410	Z472	Z534	Z596	Z658	Z720	Z782	Z844	Z906
AM	Z39	Z101	Z163	Z225	Z287	Z349	Z411	Z473	Z535	Z597	Z659	Z721	Z783	Z845	Z907
AN	Z40	Z102	Z164	Z226	Z288	Z350	Z412	Z474	Z536	Z598	Z660	Z722	Z784	Z846	Z908
AO	Z41	Z103	Z165	Z227	Z289	Z351	Z413	Z475	Z537	Z599	Z661	Z723	Z785	Z847	Z909
AP	Z42	Z104	Z166	Z228	Z290	Z352	Z414	Z476	Z538	Z600	Z662	Z724	Z786	Z848	Z910
AW	Z43	Z105	Z167	Z229	Z291	Z353	Z415	Z477	Z539	Z601	Z663	Z725	Z787	Z849	Z911
AR	Z44	Z106	Z168	Z230	Z292	Z354	Z416	Z478	Z540	Z602	Z664	Z726	Z788	Z850	Z912
AS	Z45	Z107	Z169	Z231	Z293	Z355	Z417	Z479	Z541	Z603	Z665	Z727	Z789	Z851	Z913
AT	Z46	Z108	Z170	Z232	Z294	Z356	Z418	Z480	Z542	Z604	Z666	Z728	Z790	Z852	Z914
AU	Z47	Z109	Z171	Z233	Z295	Z357	Z419	Z481	Z543	Z605	Z667	Z729	Z791	Z853	Z915
AV	Z48	Z110	Z172	Z234	Z296	Z358	Z420	Z482	Z544	Z606	Z668	Z730	Z792	Z854	Z916
AW	Z49	Z111	Z173	Z235	Z297	Z359	Z421	Z483	Z545	Z607	Z669	Z731	Z793	Z855	Z917
AX	Z50	Z112	Z174	Z236	Z298	Z360	Z422	Z484	Z546	Z608	Z670	Z732	Z794	Z856	Z918
AY	Z51	Z113	Z175	Z237	Z299	Z361	Z423	Z485	Z547	Z609	Z671	Z733	Z795	Z857	Z919
AZ	Z52	Z114	Z176	Z238	Z300	Z362	Z424	Z486	Z548	Z610	Z672	Z734	Z796	Z858	Z920

Table F (cont)


BA	Z53	Z115	Z177	Z239	Z301	Z363	Z425	Z487	Z549	Z611	Z673	Z735	Z797	Z859	Z921
BB	Z54	Z116	Z178	Z240	Z302	Z364	Z426	Z488	Z550	Z612	Z674	Z736	Z798	Z860	Z922
BC	Z55	Z117	Z179	Z241	Z303	Z365	Z427	Z489	Z551	Z613	Z675	Z737	Z799	Z861	Z923
BD	Z56	Z118	Z180	Z242	Z304	Z366	Z428	Z490	Z552	Z614	Z676	Z738	Z800	Z862	Z924
BE	Z57	Z119	Z181	Z243	Z305	Z367	Z429	Z491	Z553	Z615	Z677	Z739	Z801	Z863	Z925
BF	Z58	Z120	Z182	Z244	Z306	Z368	Z430	Z492	Z554	Z616	Z678	Z740	Z802	Z864	Z926
BG	Z59	Z121	Z183	Z245	Z307	Z369	Z431	Z493	Z555	Z617	Z679	Z741	Z803	Z865	Z927
BH	Z60	Z122	Z184	Z246	Z308	Z370	Z432	Z494	Z556	Z618	Z680	Z742	Z804	Z866	Z928
BI	Z61	Z123	Z185	Z247	Z309	Z371	Z433	Z495	Z557	Z619	Z681	Z743	Z805	Z867	Z929
BJ	Z62	Z124	Z186	Z248	Z310	Z372	Z434	Z496	Z558	Z620	Z682	Z744	Z806	Z868	Z930

In each of the above-listed aspects, the compounds include optical isomers, diastereomers, enantiomers and pharmaceutically acceptable salts thereof in isolation or mixture, where, independently at each location, the substituents R^1 , R^2 etc. are as defined herein.

- 5 Thus, for example, in one embodiment, the present invention provides a compound, or a mixture including a compound, wherein the stereochemistry of the R^1 and $C(=O)R^2$ groups are as shown in formula Ia, with R^1 and $C(=O)R^2$ in a *cis* arrangement, both over the benzo ring substituted with $-OR^6$




- 10 In another exemplary embodiment, the present invention provides a compound, or a mixture including a compound, wherein the stereochemistry of the R^1 and $C(=O)R^2$ groups are as shown in formula Ib, with R^1 and $C(=O)R^2$ in a *trans* arrangement, with only $C(=O)R^2$ over the benzo ring substituted with $-OR^6$




In yet another exemplary embodiment, the present invention provides a compound, or a mixture including a compound, having the stereochemistry of the R^1 and

C(=O)R² groups as shown in formula Ic, with R¹ and C(=O)R² in a *trans* arrangement, with only R¹ over the benzo ring substituted with -OR⁶




(Ic).

Another exemplary embodiment of the present invention provides a benzobicyclooctane compound, or a mixture containing a benzobicyclooctane compound, wherein the stereochemistry of the R¹ and C(=O)R² groups are as shown in formula Id, with R¹ and C(=O)R² in a *cis* arrangement, with neither of the R¹ nor C(=O)R² groups being over the benzo ring substituted with -OR⁶



(Id).

In one embodiment, the present invention provides a compound of formula (I) wherein R¹ is selected from the following four formulae, *i.e.*, R¹ is R1AX:



In one embodiment, the present invention provides a compound of formula (I) wherein R^1 is R1AX; R^8 is selected from hydrogen and C_1 - C_{15} alkyl; and R^9 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, (heteroaryl) C_1 - C_{15} alkylene, (C_6 - C_{10} aryl) C_1 - C_{15} alkylene, C_6 - C_{10} aryl fused to C_1 - C_{15} alkylene, (alkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, (C_6 - C_{10} aryl)(C_6 - C_{10} arylene) C_1 - C_{15} alkylene, (C_1 - C_{15} alkyl) $_p$ (heteroarylene) C_1 - C_{15} alkylene, and (heteroalkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, or two R^9 groups bonded to a common nitrogen of R^1 may be joined together to form a 5-8 membered heterocycle including the common nitrogen, where this 5-8 membered heterocycle may be substituted with 0-5 groups selected from alkyl and heteralkyl, where p is selected from 1, 2, 3, 4 and 5.

In one embodiment, the present invention provides a compound of formula I wherein R^1 is R1A and R^8 and R^9 are each independently selected from R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from (R^{11}) $_p$ -alkylene, (R^{11}) $_p$ -heteroalkylene, (R^{11}) $_p$ -arylene and (R^{11}) $_p$ -heteroarylene; R^{13} is selected from (R^{12}) $_p$ -alkylene, (R^{12}) $_p$ -heteroalkylene, (R^{12}) $_p$ -arylene, and (R^{12}) $_p$ -heteroarylene; R^{14} is selected from (R^{13}) $_p$ -alkylene, (R^{13}) $_p$ -heteroalkylene, (R^{13}) $_p$ -arylene, and (R^{13}) $_p$ -heteroarylene, R^{15} is selected from (R^{14}) $_p$ -alkylene, (R^{14}) $_p$ -heteroalkylene, (R^{14}) $_p$ -arylene, and (R^{14}) $_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. Optionally, R^9 is selected from heteroalkyl, C_1 - C_{15} alkyl, (C_6 - C_{10} aryl) C_1 - C_{15} alkylene, (C_6 - C_{10} aryl)(C_6 - C_{10} arylene) C_1 - C_{15} alkylene, (C_1 - C_{15} alkyl) $_p$ (heteroarylene) C_1 - C_{15} alkylene, and C_6 - C_{10} aryl fused to C_1 - C_{15} alkylene.

In one embodiment, the present invention provides a compound of formula (I) wherein R^1 is R1C and R^8 and R^9 are each independently selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. Optionally, R^8 is selected from hydrogen and C_1 - C_{15} alkyl; and R^9 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, (heteroaryl) C_1 - C_{15} alkylene, and (heteroalkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, or the two R^9 groups joined to a common nitrogen of R^1 may be joined together to form a 5-8 membered heterocycle including the common nitrogen.

In one embodiment, the present invention provides a compound of formula (I) wherein R^1 is R1E and R^9 is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. Optionally, R^9 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, (heteroaryl) C_1 - C_{15} alkylene, and (heteroalkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene.

In one embodiment, the present invention provides a compound of formula (I) wherein R^1 is R1F and R^9 is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene.

heteroarylene, R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. Optionally, R^9 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, (heteroaryl) C_1 - C_{15} alkylene, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, (alkyl) p (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, or the two R^9 groups of R^1 may be joined together to form a 5-8 membered heterocycle including the common nitrogen, where this 5-8 membered heterocycle may be substituted with 0-5 groups selected from alkyl and heteroalkyl.

In one embodiment, the present invention provides a compound of formula (I) wherein R^2 is $-OR^9$, *i.e.*, aspects Y1 through Y937. Optionally, R^9 of $-OR^9$ of R^2 is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene, R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. In a further optional embodiment, R^9 of $-OR^9$ of R^2 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, $(C_6$ - C_{10} aryl)(C_6 - C_{10} arylene) C_1 - C_{15} alkylene, $(C_1$ - C_{15} alkyl) p (heteroarylene) C_1 - C_{15} alkylene, $(C_1$ - C_{15} alkyl) p (heteroarylene)heteroalkylene, (heteroalkyl) p (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, and $(C_1$ - C_{15} alkyl) p (C_6 - C_{10} arylene)heteroalkylene. In a further embodiment, R^9 of $-OR^9$ of R^2 is selected from a heteroalkyl group having 1-10 carbons and 1-4 heteroatoms selected from nitrogen, oxygen, silicon and sulfur, where $-CH_2CH_2Si(CH_3)_3$ is a preferred heteroalkyl within this group.

In one embodiment, the present invention provides a compound of formula (I) wherein R^2 is $-NR^9R^9$. Optionally, R^9 of $-NR^9R^9$ of R^2 is independently selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene, R^{15} is selected from $(R^{14})_p$ -

alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. In a further optional embodiment, R^9 of $-NR^9R^9$ of R^2 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, (heteroaryl) C_1 - C_{15} alkylene, (heteroalkyl) $_p$ (aryl)heteroalkylene, (heteroalkyl) $_p$ (aryl) C_1 - C_{15} alkylene, and $(C_1$ -
5 C_{15} alkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene.

In one embodiment, the present invention provides a compound of formula (I) wherein R^3 is hydrogen.

In two embodiments, the present invention provides a compound of formula (I) wherein R^4 and R^5 are independently selected from: hydrogen, $-R^9$, $-OR^9$, and $-NR^9R^9$,
10 or R^4 and R^5 may together with the carbon to which they are both attached form a spiro carbocyclic or heterocyclic ring; and wherein R^4 and R^5 are each hydrogen. In one embodiment, the present invention provides a compound of formula (I) wherein at least one of R^4 and R^5 is selected from C_1 - C_{15} alkyl, heteroalkyl, and C_6 - C_{10} aryl. In one embodiment, the present invention provides a compound of formula (I) wherein one of R^4
15 and R^5 is hydrogen and the other of R^4 and R^5 is selected from hydrogen, $-OR^9$, $-NR^9R^9$ and $-N=N-R^9$ where the R^9 is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14}
20 is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. Optionally, R^9 of $-OR^9$, $-NR^9R^9$ and $-N=N-R^9$ from R^4 and R^5 is selected from hydrogen, C_6 - C_{10} aryl, heteroalkyl, C_1 - C_{15} alkyl, and $(C_1$ - C_{15} alkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene. In one embodiment, the
25 present invention provides a compound of formula (I) wherein R^4 and R^5 together with the carbon to which they are both attached form a 3-6-membered spiro carbocyclic or heterocyclic ring. In one embodiment, the present invention provides a compound of formula (I) wherein R^4 and R^5 together form $=O$. In one embodiment, the present invention provides a compound of formula (I) wherein R^4 and R^5 together form $=NR^{10}$ and R^{10} is $-OR^9$

where R^9 is selected from hydrogen, C_6 - C_{10} aryl, C_1 - C_8 alkyl, heteroalkyl, (C_6 - C_{10} aryl)heteroalkyl, (C_6 - C_{10} aryl) C_1 - C_{15} alkylene, (heteroalkyl) $_p$ (heteroarylene) C_1 - C_{15} alkylene, (heteroalkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, and (C_1 - C_{15} alkyl) $_p$ (C_6 - C_{10} arylene)heteroalkylene. In one embodiment, the present invention provides a compound of formula (I) wherein R^4 and R^5 together form $=NR^{10}$ and R^{10} is $-N(R^9)(R^9)$ where R^9 is selected from hydrogen, C_1 - C_8 alkyl, heteroalkyl, C_6 - C_{10} aryl, (C_6 - C_{10} aryl)heteroalkylene, (heteroalkyl) $_p$ C_6 - C_{10} arylene, (C_1 - C_{15} alkyl) $_p$ C_6 - C_{10} arylene, (heteroalkyl) $_p$ (C_6 - C_{10} arylene)heteroalkylene, (C_1 - C_{15} alkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, and (C_1 - C_{15} alkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} heteroalkylene. In one embodiment, the present invention provides a compound of formula (I) wherein R^4 and R^5 together form $=CR^8R^8$, and one of R^8 is hydrogen while the other R^8 is selected from hydrogen, C_1 - C_8 alkyl and heteroalkyl.

In one embodiment, the present invention provides a compound of formula (I) wherein R^6 is hydrogen. In another embodiment, the present invention provides a compound of formula (I) wherein R^6 is selected from R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from (R^{11}) $_p$ -alkylene, (R^{11}) $_p$ -heteroalkylene, (R^{11}) $_p$ -arylene and (R^{11}) $_p$ -heteroarylene; R^{13} is selected from (R^{12}) $_p$ -alkylene, (R^{12}) $_p$ -heteroalkylene, (R^{12}) $_p$ -arylene, and (R^{12}) $_p$ -heteroarylene; R^{14} is selected from (R^{13}) $_p$ -alkylene, (R^{13}) $_p$ -heteroalkylene, (R^{13}) $_p$ -arylene, and (R^{13}) $_p$ -heteroarylene; R^{15} is selected from (R^{14}) $_p$ -alkylene, (R^{14}) $_p$ -heteroalkylene, (R^{14}) $_p$ -arylene, and (R^{14}) $_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. In another embodiment, R^6 is selected from C_1 - C_{15} alkyl, C_1 - C_{15} heteroalkyl, (C_6 - C_{10} aryl) C_1 - C_{15} alkylene, (C_6 aryl)(C_6 aryl) C_1 - C_{15} alkylene, (C_2 - C_6 heteroaryl) C_1 - C_{15} alkylene, (C_6 - C_{10} aryl) C_1 - C_{15} heteroalkylene, (heteroalkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene, (heteroalkyl) $_p$ (C_2 - C_6 heteroarylene) C_1 - C_{15} alkylene, and (heteroalkyl) $_p$ (C_6 arylene)(heteroalkylene)(C_6 arylene) C_1 - C_{15} alkylene. In one embodiment of the present invention, R^6 is as defined above with the proviso that R^6 is not lower alkyl, *e.g.*, is not C_1 - C_6 so that $-OR^6$ is not C_1 - C_6 alkoxy.

In one embodiment, the present invention provides a compound of formula (I) wherein R^8 is hydrogen.


In one embodiment, the present invention provides a compound of formula (I) wherein n is 0. In another embodiment, the present invention provides a compound of formula (I) wherein n is 1. In another embodiment, the present invention provides a compound of formula (I) wherein n is 0 or 1.

5 In one embodiment, the present invention provides a compound of formula (I) wherein $-R^1$ is *trans* to $-C(O)R^2$, i.e., compounds of formula (Ib) and (Ic), also referred to herein as compounds of formula (Ih).

In one embodiment, the present invention provides a compound of formula (I) wherein $-R^1$ is *cis* to $-C(O)R^2$, i.e., compounds of formula (Ia) and (Id), also referred to
10 herein as compounds of formula (Ig).


In one embodiment, the present invention provides a compound of formula (I) wherein R^3 is hydrogen; R^4 and R^5 are selected from (a) R^4 is hydrogen and R^5 is hydroxyl or protected hydroxyl and (b) R^4 and R^5 together form carbonyl; R^6 is hydrogen; and n is 0. In one embodiment R^2 is $-OR^9$ where a preferred R^2 group is
15 $-OCH_2CH_2Si(CH_3)_3$.

In one embodiment R^1 is



where optionally R^9 is a C_1 - C_6 hydrocarbonyl, such as, in one embodiment, n-propyl and $-\text{CH}_2\text{-CH=CH}_2$.

In one embodiment R^1 is




20 where optionally R^8 is hydrogen and R^9 is C_1 - C_6 hydrocarbonyl, such as, in one embodiment, R^9 is $-\text{CH}_2\text{-CH=CH}_2$.

B. Preparation of Benzobicyclooctane Compounds

The benzobicyclooctanes of this invention may be prepared according to Schemes 1-4. In these Schemes, "PG" denotes a protecting group. Suitable protecting groups are set forth in, for example, Greene and Wuts, *Protective Groups in Organic Synthesis*, 2d Edition, John Wiley & Sons, New York, 1991.

Scheme 1



In Scheme 1, the starting material (not shown) for 1 may be prepared by the Diels-Alder reaction of 2,7-dihydroxynaphthalene with maleic anhydride (*see, e.g.,* Singh, A.K.; Yadar, S.; Bhattacharjee, G., *J. Indian Chemical Soc.* 1990, 67, 818; and Takeda, K.; Hagishita, S.; Sugiura, M.; Kitahonoki, K.; Ban, I.; Miyazaki, S.; Kuriyama, K., *Tetrahedron* 1970, 26, p. 1435). The resulting anhydride may be opened with a suitable alcohol, *e.g.,* trimethylsilylethanol, to give the 9-protected and the 10-protected benzobicyclooctane, 1 (only the 9-ester is depicted).

In Scheme 1, chemical steps a, b, c, d, and e represent the following reaction conditions.

(a) is a chemical reaction wherein the free acid of 1 is transformed into the reactive intermediate 2. Suitable conditions for this type of reaction involve treating 1

with a suitable activating agent, *e.g.*, diphenylphosphoryl azide, in the presence of a suitable base, *e.g.*, an organoamine such as diisopropylethylamine (DIEA), in an appropriate solvent, *e.g.*, tetrahydrofuran (THF), at a suitable reaction temperature, *e.g.*, at ambient temperature. Alternatively, formation of an active ester via a suitable coupling agent and hydroxy compound, *e.g.*, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI) and 1-hydroxybenzotriazole (HOBt), under the same conditions produce **2**, suitable for use in steps b or c. In either process, X is a leaving group that activates the adjoining carbonyl group.

(b) is a chemical reaction in which the activated acid **2** forms the ester **3**. Suitable conditions for this type of reaction involve treating **2** with a suitable alcohol (R^9OH), *e.g.*, n-propanol, in the presence of a suitable catalyst, *e.g.*, 4-dimethylaminopyridine (DMAP), in an appropriate solvent, *e.g.*, THF, at an appropriate temperature, *e.g.*, ambient temperature. In alcohols of formula R^9OH , R^9 is an organic group having 1-30 carbons and optionally containing 1-4 heteroatoms selected from nitrogen, oxygen, silicon and sulfur, with the provision that two R^9 groups both joined to a common atom may be joined together so as to form a ring with the common atom. In one embodiment, R^9 is selected from R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5. Optionally, R^9 is selected from heteroalkyl, C_1 - C_{15} alkyl, $(C_6$ - C_{10} aryl) C_1 - C_{15} alkylene, $(C_6$ - C_{10} aryl)(C_6 - C_{10} arylene) C_1 - C_{15} alkylene, $(C_1$ - C_{15} alkyl) $_p$ (heteroarylene) C_1 - C_{15} alkylene, and C_6 - C_{10} aryl fused to C_1 - C_{15} alkylene. Numerous suitable alcohols of formula R^9OH are either commercially available chemicals or are compounds described in the chemical literature.


(c) is a chemical reaction in which **2** is coupled with an amine to give the amide **4**. Suitable conditions for this type of reaction involve treating **2** with a suitable

amine (R^9R^9NH), *e.g.*, di(*n*-pentyl)amine, and a suitable base (if required), *e.g.*, DIEA, in the presence of a suitable catalyst, *e.g.*, DMAP, in an appropriate solvent, *e.g.*, THF, at ambient temperature. In amines of formula R^9R^9NH , R^9 is selected from hydrogen and organic groups having 1-30 carbons and optionally containing 1-4 heteroatoms selected from nitrogen, oxygen, silicon and sulfur, with the proviso that the two R^9 groups may be joined together so as to form a ring with the nitrogen to which they are both attached. In one embodiment, R^9 is selected from hydrogen, R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene; and p is selected from 0, 1, 2, 3, 4 and 5. Optionally, R^9 is selected from hydrogen, heteroalkyl, C_1 - C_{15} alkyl, (heteroaryl) C_1 - C_{15} alkylene, and (heteroalkyl) $_p$ (C_6 - C_{10} arylene) C_1 - C_{15} alkylene. Numerous suitable amines of formula R^9R^9NH are either commercially available chemicals or are compounds described in the chemical literature.

(d) is a chemical reaction in which **2** is an acyl azide and is converted to the corresponding isocyanate prior to reaction with an alcohol (R^9OH as defined above) to yield carbamate **5**. Suitable conditions for this type of reaction involve first heating **2** in suitable solvent, *e.g.*, refluxing dioxane, and then treating the resulting isocyanate with a suitable alcohol R^9OH , *e.g.*, *n*-propanol, in the absence or presence of a suitable catalyst, *e.g.*, DMAP.

(e) is a chemical reaction in which **2** is an acyl azide and is converted to the isocyanate prior to reacting with an amine (R^9R^9NH as defined above), to yield urea **6**. Suitable conditions for this type of reaction involve first heating **2** in a suitable solvent, *e.g.*, refluxing dioxane, and then treating the resulting isocyanate with a suitable amine (R^9R^9NH), *e.g.*, morpholine or tyramine.

Scheme 2



In Scheme 2, chemical steps f, g and h represent the following reaction conditions.


(f) is a chemical reaction in which the protecting group of **3** is removed to give **7**. When, for example, PG is trimethylsilylethyl, it may be removed by exposure to a suitable fluoride source, *e.g.*, tetrabutylammonium fluoride (TBAF), in a suitable solvent, *e.g.*, anhydrous THF. Alternatively, suitable deprotection conditions involve performing an acidolysis in, *e.g.*, TFA/H₂O, 9/1 (v/v). Other conditions for removing protecting groups are set forth in Greene and Wuts, *Protective Groups in Organic Synthesis*, 2d Edition, John Wiley & Sons, New York, 1991.

(g) is a chemical reaction in which **7** is coupled to an alcohol to give **8**. Suitable conditions for this type of reaction involve treating **7** with a suitable alcohol, *e.g.*, dimethylbutanol, a coupling reagent such as O-(N-Succinimidyl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TSTU), a suitable base, *e.g.*, an organoamine such as N-methylmorpholine (NMM), in the presence of a suitable catalyst, *e.g.*, DMAP, in an appropriate solvent, *e.g.*, 5% dimethylformamide (DMF) in THF.

(h) is a chemical reaction in which **7** is coupled with an amine HNR⁹R⁹ to give **9**. Suitable conditions for this type of reaction involve treating **7** with a suitable coupling reagent such as O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium

hexafluorophosphate (HATU), a suitable base, *e.g.*, an organoamine such as NMM, in the presence of a suitable catalyst, *e.g.*, DMAP, in an appropriate solvent, *e.g.*, THF.

Scheme 3



5

In Scheme 3, chemical steps i, j, k, l and m represent the following reaction conditions.

(i) is a chemical reaction in which the ketone group of **3** is derivatized with an organohydrazine or organohydroxylamine to give **10**. Suitable conditions for performing this type of reaction involve treating the ketone with a suitable hydrazine or hydroxylamine, *e.g.*, methyl hydrazine or O-phenyl-hydroxylamine, in a suitable solvent, *e.g.*, methanol.

(j) is a chemical reaction in which the ketone group of **3** is reduced to give alcohols **11** and **12**. Suitable conditions for performing this type of reaction involve treating the ketone with a suitable reducing agent, *e.g.*, NaBH₄, in a suitable solvent, *e.g.*, methanol. Other suitable reducing conditions are set forth in well known books and

treatises. The resulting stereoisomers **11** and **12** may be separated from one another by, *e.g.*, column chromatography.


(k) is a chemical reaction in which the ketone of group of **3** undergoes reductive amination to give amines **13** and **14**. Suitable conditions for performing this type of reaction involve treating the ketone with a suitable amine (HNR^9R^9), *e.g.*, dimethylamine, a suitable reducing agent, *e.g.*, NaBH_3CN , in the presence of a mild acid, *e.g.*, acetic acid, in a suitable solvent, *e.g.*, methanol. Other suitable reductive amination conditions are set forth in well known books and treatises. The resulting stereoisomers **13** and **14** may be separated from one another by, *e.g.*, column chromatography.

10 (l) is a chemical reaction in which the phenolic group of **3** is alkylated to give **15**. Suitable conditions for performing this type of reaction involve treating **3** with a suitable alkyl halide, *e.g.*, N,N-diethyl-2-chloroacetamide, in the presence of a suitable inorganic base, *e.g.*, Cs_2CO_3 , in a suitable solvent, *e.g.*, dimethoxyethane (DME) or DMF. Other suitable alkyl halides of formula $\text{R}^6\text{-X}$ are well known in the art, where X is halide, and R^6 is selected from R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(\text{R}^{11})_p\text{-alkylene}$, $(\text{R}^{11})_p\text{-heteroalkylene}$, $(\text{R}^{11})_p\text{-arylene}$ and $(\text{R}^{11})_p\text{-heteroarylene}$; R^{13} is selected from $(\text{R}^{12})_p\text{-alkylene}$, $(\text{R}^{12})_p\text{-heteroalkylene}$, $(\text{R}^{12})_p\text{-arylene}$, and $(\text{R}^{12})_p\text{-heteroarylene}$; R^{14} is selected from $(\text{R}^{13})_p\text{-alkylene}$, $(\text{R}^{13})_p\text{-heteroalkylene}$, $(\text{R}^{13})_p\text{-arylene}$, and $(\text{R}^{13})_p\text{-heteroarylene}$, R^{15} is selected from $(\text{R}^{14})_p\text{-alkylene}$, $(\text{R}^{14})_p\text{-heteroalkylene}$, $(\text{R}^{14})_p\text{-arylene}$, and $(\text{R}^{14})_p\text{-heteroarylene}$, and p is selected from 0, 1, 2, 3, 4 and 5, and optionally is selected from $\text{C}_1\text{-C}_{15}\text{alkyl}$, $\text{C}_1\text{-C}_{15}\text{heteroalkyl}$, $(\text{C}_6\text{-C}_{10}\text{aryl})\text{C}_1\text{-C}_{15}\text{alkylene}$, $(\text{C}_6\text{aryl})(\text{C}_6\text{aryl})\text{C}_1\text{-C}_{15}\text{alkylene}$, $(\text{C}_2\text{-C}_6\text{heteroaryl})\text{C}_1\text{-C}_{15}\text{alkylene}$, $(\text{C}_6\text{-C}_{10}\text{aryl})\text{C}_1\text{-C}_{15}\text{heteroalkylene}$, $(\text{heteroalkyl})_p(\text{C}_6\text{-C}_{10}\text{arylene})\text{C}_1\text{-C}_{15}\text{alkylene}$, $(\text{heteroalkyl})_p(\text{C}_2\text{-C}_6\text{heteroarylene})\text{C}_1\text{-C}_{15}\text{alkylene}$, and $(\text{heteroalkyl})_p(\text{C}_6\text{arylene})(\text{heteroalkylene})(\text{C}_6\text{arylene})\text{C}_1\text{-C}_{15}\text{alkylene}$. Numerous suitable alkyl halides are either commercially available chemicals or are compounds described in the chemical literature.

(m) is a chemical reaction in which the phenolic group of **3** is alkylated to give **15**. Suitable conditions for performing this type of reaction involve treating **3** with

an organic compound having a suitably activated hydroxyl group in a suitable solvent, such as THF. For example, allyl 4-hydroxymethylbenzoate may be activated by exposure to a phosphine, *e.g.*, triphenylphosphine, and a suitable azo compound, *e.g.*, diethylazodicarboxylate (DEAD). Other suitable compounds having an activated hydroxyl group may be readily prepared from the corresponding alcohol of the formula R^6-OH where R^6 is an organic group. Alcohols of the formula R^6-OH are well known in the art, including alcohols wherein R^6 is selected from R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene, $(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene; R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5, and optionally is selected from C_1 - C_{15} alkyl, C_1 - C_{15} heteroalkyl, $(C_6-C_{10}aryl)C_1$ - C_{15} alkylene, $(C_6aryl)(C_6aryl)C_1$ - C_{15} alkylene, $(C_2-C_6heteroaryl)C_1$ - C_{15} alkylene, $(C_6-C_{10}aryl)C_1$ - C_{15} heteroalkylene, $(heteroalkyl)_p(C_6-C_{10}arylene)C_1$ - C_{15} alkylene, $(heteroalkyl)_p(C_2-C_6heteroarylene)C_1$ - C_{15} alkylene, and $(heteroalkyl)_p(C_6arylene)(heteroalkylene)(C_6arylene)C_1$ - C_{15} alkylene. Numerous suitable alcohols are either commercially available chemicals or are compounds described in the chemical literature.

Scheme 4



In Scheme 4, chemical steps n, o, p and q represent the following reaction conditions.

(n) is a chemical reaction wherein the ester-carbamate **5** (prepared in, *e.g.*, Scheme 1) is transformed into the corresponding ester-amine **16**. Suitable conditions for this type of reaction involve treating **5** under reducing conditions, *e.g.*, H_2 , on a suitable catalyst or solid support, *e.g.*, palladium, in the presence of a suitable solvent, *e.g.*, ethanol.

(o) is a chemical reaction wherein the ester-amine **16** is acylated to form the corresponding ester-amide **17**. Suitable conditions for this type of reaction involve treating **16** with an acylating agent, generally denoted as $R^9-C(=O)-X$ where R^9 represents R^9 as set forth in compounds of formula **R1a**, and X is a leaving group, *e.g.*, chloride. The acylation reaction is suitably conducted in the presence of an amine, such as a secondary or tertiary amine, *e.g.*, diisopropylethylamine (DIEA).

(p) is a chemical reaction wherein the ester-1°amine **16** is transformed into an ester-2°amine **18**. Suitable conditions for this type of reaction involve treating **16** with an aldehyde of the formula R^8-CHO , in the presence of a reducing agent, *e.g.*, $NaCNBH_3$. In Scheme 4, the designation " R^8 " is used to denote the " R^8 " group as found in, for example, compound of formula **R1a**. Compounds of formula R^8-CHO wherein R^8 is selected from alkyl, aryl and heteroalkyl are well known in the chemical literature, and are available from commercial suppliers of chemicals. The ester-2°amine **18** is a suitable intermediate in the preparation of compounds of formula **19**, which are also compounds of formula **R1a**.

(q) is a chemical reaction wherein the ester-2°amine **18** is transformed into an ester-amide **19**. Suitable conditions for this type of reaction involve treating **18** with an acylating agent, generally denoted as $R^9-C(=O)-X$, where R^9 is used in Scheme 4 to denote " R^9 " in, for example, compounds of formula **R1a**, and X is a leaving group, *e.g.*, chloride. The acylation reaction is suitably conducted in the presence of an amine, such as a secondary or tertiary amine, *e.g.*, diisopropylethylamine (DIEA). Compounds of the formula $R^9-C(=O)-X$ are readily prepared from the corresponding carboxylic acid of the formula $R^9-C(=O)-OH$ by treatment with, *e.g.*, thionyl chloride.

Numerous compounds of the formula $R^9-C(=O)-OH$ wherein R^9 is an organic group having 1-30 carbons and optionally containing 1-4 heteroatoms selected from nitrogen, oxygen and silicon are well known in the chemical literature, and/or may be obtained from many commercial suppliers of chemicals. Furthermore, many compounds of formula $R^9-C(=O)-OH$ wherein R^9 is selected from R^{11} , R^{12} , R^{13} , R^{14} and R^{15} where R^{11} is selected from alkyl, heteroalkyl, aryl and heteroaryl; R^{12} is selected from $(R^{11})_p$ -alkylene,

$(R^{11})_p$ -heteroalkylene, $(R^{11})_p$ -arylene and $(R^{11})_p$ -heteroarylene; R^{13} is selected from $(R^{12})_p$ -alkylene, $(R^{12})_p$ -heteroalkylene, $(R^{12})_p$ -arylene, and $(R^{12})_p$ -heteroarylene; R^{14} is selected from $(R^{13})_p$ -alkylene, $(R^{13})_p$ -heteroalkylene, $(R^{13})_p$ -arylene, and $(R^{13})_p$ -heteroarylene, R^{15} is selected from $(R^{14})_p$ -alkylene, $(R^{14})_p$ -heteroalkylene, $(R^{14})_p$ -arylene, and $(R^{14})_p$ -heteroarylene, and p is selected from 0, 1, 2, 3, 4 and 5, are well known in the chemical literature and/or may be obtained from commercial suppliers of chemicals. Furthermore, many compounds of formula $R^9-C(=O)-OH$ wherein R^9 is selected from heteroalkyl, C_1-C_{15} alkyl, (heteroaryl) C_1-C_{15} alkylene, $(C_6-C_{10}aryl)C_1-C_{15}$ alkylene, $C_6-C_{10}aryl$ fused to C_1-C_{15} alkylene, $(alkyl)_p(C_6-C_{10}arylene)C_1-C_{15}$ alkylene, $(C_6-C_{10}aryl)(C_6-C_{10}arylene)C_1-C_{15}$ alkylene, $(C_1-C_{15}alkyl)_p(heteroarylene)C_1-C_{15}$ alkylene, and $(heteroalkyl)_p(C_6-C_{10}arylene)C_1-C_{15}$ alkylene, are well known in the chemical literature and/or may be obtained from commercial suppliers of chemicals. Furthermore, many compounds of formula $R^9-C(=O)-OH$ wherein R^9 is selected from heteroalkyl, C_1-C_{15} alkyl, $(C_6-C_{10}aryl)C_1-C_{15}$ alkylene, (heteroaryl) C_1-C_{15} alkylene, and $(heteroalkyl)_p(C_6-C_{10}arylene)C_1-C_{15}$ alkylene, are well known in the chemical literature and/or may be obtained from commercial suppliers of chemicals. Furthermore, many compounds of formula $R^9-C(=O)-OH$ wherein R^9 is selected from heteroalkyl, C_1-C_{15} alkyl, $(C_6-C_{10}aryl)C_1-C_{15}$ alkylene, $(C_6-C_{10}aryl)(C_6-C_{10}arylene)C_1-C_{15}$ alkylene, $(C_1-C_{15}alkyl)_p(heteroarylene)C_1-C_{15}$ alkylene, and $C_6-C_{10}aryl$ fused to C_1-C_{15} alkylene are well known in the chemical literature and/or may be obtained from commercial suppliers of chemicals. Furthermore, many compounds of formula $R^9-C(=O)-OH$ wherein R^9 is selected from heteroalkyl, C_1-C_{15} alkyl, (heteroaryl) C_1-C_{15} alkylene, and $(heteroalkyl)_p(C_6-C_{10}arylene)C_1-C_{15}$ alkylene are well known in the chemical literature and/or may be obtained from commercial suppliers of chemicals. Furthermore, many compounds of formula $R^9-C(=O)-OH$ wherein R^9 is selected from heteroalkyl, C_1-C_{15} alkyl, (heteroaryl) C_1-C_{15} alkylene, $(C_6-C_{10}aryl)C_1-C_{15}$ alkylene, $(C_6-C_{10}aryl)(C_6-C_{10}arylene)C_1-C_{15}$ alkylene, $(C_1-C_{15}alkyl)_p(heteroarylene)C_1-C_{15}$ alkylene, and $C_6-C_{10}aryl$ fused to C_1-C_{15} alkylene are well known in the chemical literature and/or may be obtained from commercial suppliers of chemicals.

C₁₀aryl)C₁-C₁₅alkylene, (alkyl)p(C₆-C₁₀arylene)C₁-C₁₅alkylene, are well known in the chemical literature and/or may be obtained from commercial suppliers of chemicals. These carboxylic acids may be used in the preparation of compounds of the present invention.


One skilled in the art of organic synthesis would readily understand that the chemical steps disclosed above may be performed in a variety of sequences to produce bicyclooctanes of this invention. For instance, the compound of Example 2 undergoes step (h) to give the compound of Example 4. This compound in turn undergoes step (m) to give the bicyclooctane of Example 5.

The present invention provides benzobicyclooctane compounds wherein R³ may or may not be hydrogen, and independently, R⁷ may replace a hydrogen either 0, 1, 2 or 3 times on the "benzo" portion of the benzobicyclooctane compound. Compounds wherein R³ is hydrogen and n is 0 are readily prepared from (unsubstituted) 2,7-dihydroxynaphthalene, as shown in Schemes 1, 2, 3 and 4. Compounds wherein R³ is not hydrogen, and/or n is not 0, are readily prepared from the corresponding substituted 2,7-dihydroxynaphthalene. For example, a benzobicyclooctane compound of the invention wherein R³ is methyl and n is 1 with R⁷ being a methyl group may be prepared from a dimethyl substituted 2,7-dihydroxynaphthalene, e.g., 2,7-dihydroxy-3,6-dimethylnaphthalene as shown in Scheme 5. Commercial supply houses, custom chemical supply houses, and published synthetic methods provide access to a large number of substituted 2,7-dihydroxynaphthalene compounds that may be used in preparing compounds of the present invention.

Furthermore, benzobicyclooctane compounds wherein R³ is not equal to hydrogen and/or n is 1, 2 or 3 may be used in the synthetic transformations shown in Schemes 1, 2, 3 and/or 4, in lieu of the hydrogen-substituted benzobicyclooctane depicted in those Schemes, to provide compounds of the present invention. For instance, the benzobicyclooctane produced by the Diels-Alder reaction of maleic anhydride and 2,7-dihydroxy-3,6-dimethylnaphthalene as shown in Scheme 5 may be treated to open up the anhydride and form the corresponding acid/ester. Exemplary treatment conditions are DMAP with trimethylsilylethanol (*see, e.g.,* Example 1 as described herein), which

provides the corresponding carboxylic acid/trimethylsilylethylene ester as shown in Scheme 5, where this acid/ester is a representative compound of formula 1 as shown in Schemes 1, 2, 3 and 4:

Scheme 5



5

In one aspect, the present invention provides benzobicyclooctane compounds wherein R^6 is hydrogen or an organic group having 1-20 carbons, wherein the organic group may optionally include 1-4 heteroatoms selected from nitrogen, oxygen, silicon and sulfur. Schemes 1, 2, 3 and 4 illustrate synthetic methodology using a benzobicyclooctane compound wherein R^6 is hydrogen. However, the same methodology may be employed with benzobicyclooctane compounds wherein R^6 is an organic group.


Alternatively, a compound of the invention may be prepared according to Schemes 1, 2, 3 and 4, having desired R^1 , R^2 , R^3 , R^4 , R^5 and R^7 groups, with R^6 being hydrogen. The R^6 hydrogen may be replaced with an organic group having 1-20 carbons

and optionally having 1-4 heteroatoms selected from nitrogen, oxygen, silicon and sulfur, as shown in steps l or m of Scheme 3. This later approach is illustrated in several Examples as set forth herein (*see, e.g.*, Examples 5, 10, 33 (describing General Procedure F for converting $R^6=H$ to $R^6=\text{organic group}$), 35 (describing General Procedure G) 36-43 and 5 45 (employing General Procedure F), and 44, 46-52 (employing General Procedure G) and 87-88. See also Examples 7, 8 and 34 wherein the R^6 group is replaced with a different R_6 group). Accordingly, in view of the present disclosure, those of ordinary skill in the art can prepare compounds of the present invention wherein R^6 is hydrogen or an organic group.

Benzobicyclooctane compounds of the invention wherein R^6 is an inorganic
10 group having 1-8 atoms exclusively selected from boron, sulfur, phosphorous, silicon and hydrogen, may readily be prepared from the corresponding phenolic compound, *i.e.*, compounds wherein R^6 is H. Methodology to convert alcohols to, *e.g.*, sulfates, sulfonates, phosphates, phosphonates, borates, and boronates, where these groups are exemplary inorganic R^6 groups, are well known in the art, and may be employed in the preparation of
15 compounds of the present invention. For clarification, it will be noted that groups including heteroatoms as well as carbon atoms, *e.g.*, $-O-B(OR)_2$ and $-S(O)_2R$ where R is an organic group, are included within the scope of heteroalkyls as defined herein.


The present invention provides various stereoisomers of benzobicyclooctanes, in isolated form or as mixtures of stereoisomers, and in particular
20 provides the diastereomers shown as Formulae Ia, Ib, Ic and Id. Any of these four diastereomers can be prepared according to the present invention. The Diels-Alder reaction of 2,7-dihydroxynaphthalene and maleic acid typically forms two diastereomers, shown as structures A and B in Scheme 6.

Scheme 6




The diastereomers **A** and **B** can be separated from one another by, for example, chromatography, and then each can be reacted individually with trimethylsilyl
5 ethanol to provide a mixture of the corresponding two *cis* acid-esters (**C** and **D**), as shown in Scheme 7a starting from diastereomer **A**, or the corresponding two *trans* acid-esters (**E** and **F**), as shown in Scheme 7b starting from diastereomer **B**.

Scheme 7a



Scheme 7b




5 The diastereomers **C** and **D** may be separated from one another by, for example, chromatography. Likewise, the diastereomers **E** and **F** can be separated from one another by, for example, chromatography. Each of the diastereomers **C**, **D**, **E** and **F** may be


reacted under conditions to give either the *trans* or *cis* products. For example, as shown in Scheme 8a, diastereomer **C** may be reacted to form the *trans* diastereomer **G** or the *cis* diastereomer **H** where X is -OR (diester) or -NRR (ester amide). Likewise, diastereomer **D** may be reacted to form *cis* and *trans* products as shown in Scheme 8b.

5

Scheme 8a




Scheme 8b



C. Libraries

In one aspect, the present invention provides a library of 5 benzobicyclooctane compounds. In one aspect the library includes, *i.e.*, comprises, a plurality of compounds each having a structure of formula (I), while in another aspect the library consists of a plurality of compounds each having a structure of formula (I)




(I)

where each of R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , and R^7 have been defined above, including narrower embodiments thereof and set forth above, and diversity is present among the R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , and R^7 groups.


A library according to the present invention may be prepared by combinatorial synthetic techniques, where such a library is referred to herein as a combinatorial library. An exemplary combinatorial approach to preparing a library of the present invention is a solid-phase technique, where the benzobicyclooctane scaffold is covalently attached to a solid support. An exemplary solid-phase combinatorial technique includes the following steps:

- 10 (a) providing a compound bound to a solid support according to formula (II)



- wherein PG1 and PG2 refer to first and second protecting groups, respectively, where the first protecting group can be removed in the continued presence of the second protecting group, and the second protecting group can be removed in the continued presence of the linker, and (SS) refers to a solid support;
- 15


- (b) removing the first protecting group but not the second protecting group, to provide a first deprotected product;
- (c) reacting the first deprotected product with a plurality of amines of the formula HNR^1R^2 to provide a plurality of compounds bound to a solid support, each according to formula (IIa)
- 20



where R and R' are each independently selected from R⁹;


(d) removing the second protecting group from (IIa) to provide a second deprotected product;

- 5 (e) reacting the second deprotected product with a plurality of amines of the formula HNR''R''' to provide a plurality of compounds bound to a solid support, each according to formula (IIb)




where R'' and R''' are each independently selected from R⁹; and

- 10 (f) removing the benzobicyclooctane compounds from the linker to provide a library of compounds according to formula (IIc)



In various embodiments of this method: PG1 is $-\text{CH}_2\text{-CH=CH}_2$; and/or wherein PG2 is $-\text{CH}_2\text{CH}_2\text{-Si(CH}_3)_3$; and/or linker is



; or PG1 is $-\text{CH}_2\text{-CH=CH}_2$ while PG2 is $-\text{CH}_2\text{CH}_2\text{-Si(CH}_3)_3$ and while linker is $\text{O}-\text{CH}_2-\text{C}_6\text{H}_4-\text{O}-\text{CH}_2-\text{C}(=\text{O})-\text{NH}-\text{[SS]}$

- 5 In various embodiments, additionally or alternatively: removing the first protecting group but not the second protecting group, to provide a first deprotected product according to step (b), is accomplished by reacting (II) with $\text{Pd(PPh}_3)_4$ and N-methylaniline; and/or removing the second protecting group from (IIa) to provide a second deprotected product according to step (d) is accomplished by treating (IIa) with tetrabutylammonium
10 fluoride solution; and/or removing the linker to provide a library of compounds according to formula (IIc) is accomplished by treating (IIb) with aqueous trifluoroacetic acid.

In various embodiments, additionally or alternatively, the library prepares compounds wherein R^3 is H, R^4 and R^5 collectively form $=\text{O}$, and n is zero.

C. Pharmaceutical Compositions

- 15 In another aspect, the present invention provides a composition containing a benzobicyclooctane compound of formula (I) in admixture with a pharmaceutically acceptable adjuvant, carrier, diluent or excipient, *i.e.*, the present invention provides a pharmaceutical composition containing a compound of formula (I). In other aspects, the present invention provides a composition containing a benzobicyclooctane compound
20 according to each of embodiments, X1-X930, Y1-Y930 and Z1-Z930 in admixture with a pharmaceutically acceptable adjuvant, carrier, diluent or excipient. The pharmaceutical composition may contain optional ingredient(s) if desired.

The pharmaceutical compositions of the present invention may be in any form which allows for the composition to be administered to a patient. Typical routes of
25 administration include, without limitation, oral, topical, parenteral, sublingual, rectal,

vaginal, and intranasal. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. Pharmaceutical composition of the invention are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a patient. Compositions that will be administered to a patient take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of benzobicyclooctane in aerosol form may hold a plurality of dosage units.

The composition may be in the form of a solid, liquid or gas (aerosol). In one aspect, the carrier(s) are particulate, so that the compositions are, for example, in tablet or powder form. The carrier(s) may be liquid, with the compositions being, for example, an oral syrup or injectable liquid. In addition, the carrier(s) may be gaseous, so as to provide an aerosol composition useful in, *e.g.*, inhalatory administration.

When intended for oral administration, the composition is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid.

As a solid composition for oral administration, the composition may be formulated into a powder, granule, compressed tablet, pill, capsule, chewing gum, wafer or the like form. Such a solid composition will typically contain one or more inert diluents or edible carriers. In addition, one or more of the following adjuvants may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch, lactose or dextrans, disintegrating agents such as alginic acid, sodium alginate, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin, a flavoring agent such as peppermint, methyl salicylate or orange flavoring, and a coloring agent.

When the composition is in the form of a capsule, *e.g.*, a gelatin capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or a fatty oil.

The composition may be in the form of a liquid, *e.g.*, an elixir, syrup, solution, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred composition contain, in addition to the present compounds, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be
5 administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

The liquid pharmaceutical compositions of the invention, whether they be solutions, suspensions or other like form, may include one or more of the following
10 adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite;
15 chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

20 A liquid compositions intended for either parenteral or oral administration should contain an amount of the inventive compound such that a suitable dosage will be obtained. Typically, this amount is at least 0.01% of a compound of the invention in the composition. When intended for oral administration, this amount may be varied to be between 0.1 and about 70% of the weight of the composition. Preferred oral compositions
25 contain between about 4% and about 50% of the active vanadium(V) complex. Preferred compositions and preparations according to the present invention are prepared so that a parenteral dosage unit contains between 0.01 to 1% by weight of active compound.

The pharmaceutical composition may be intended for topical administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base.

The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a pharmaceutical composition for topical administration. If intended for transdermal administration, the
5 composition may include a transdermal patch or iontophoresis device. Topical formulations may contain a concentration of the inventive compound of from about 0.1 to about 10% w/v (weight per unit volume).

The composition may be intended for rectal administration, in the form, *e.g.*, of a suppository which will melt in the rectum and release the drug. The composition for
10 rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter and polyethylene glycol.

The composition may include various materials which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials which form the coating shell
15 are typically inert, and may be selected from, for example, sugar, shellac, and other enteric coating agents. Alternatively, the active ingredients may be encased in a gelatin capsule.

The composition in solid or liquid form may include an agent which binds to the benzobicyclooctane compounds of the invention and thereby assists in the delivery of the active compound. Suitable agents which may act in this capacity include a
20 monoclonal or polyclonal antibody, a protein or a liposome.

Materials used in preparing the pharmaceutical compositions should be pharmaceutically pure and non-toxic in the amounts used. It will be evident to those of ordinary skill in the art that the optimal dosage of the active ingredient(s) in the pharmaceutical composition will depend on a variety of factors. Relevant factors include,
25 without limitation, the type of subject (*e.g.*, human), the particular form of the active ingredient, the manner of administration and the composition employed.

The pharmaceutical composition of the present invention may consist of gaseous dosage units, *e.g.*, it may be in the form of an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of

pressurized packages. Delivery may be by a liquefied or compressed gas or by a suitable pump system which dispenses the active ingredients. Aerosols of compounds of the invention may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, 5 activators, valves, subcontainers, and the like, which together may form a kit. Preferred aerosols may be determined by one skilled in the art, without undue experimentation.

Whether in solid, liquid or gaseous form, the pharmaceutical composition of the present invention may contain one or more known pharmacological agents used in the treatment of inflammation.

10 The pharmaceutical compositions may be prepared by methodology well known in the pharmaceutical art. For example, a composition intended to be administered by injection can be prepared by combining a benzobicyclooctane compounds of formula (I) with water so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that non-covalently 15 interact with the benzobicyclooctane compound so as to facilitate dissolution or homogeneous suspension of the compound in the aqueous delivery system.

D. Biological Applications

The present invention provides benzobicyclooctanes, compositions containing a benzobicyclooctane, and methods of using benzobicyclooctane compounds to 20 inhibit cellular events involving TNF- α or IL-8. Thus, in one aspect, the present invention provides a method to modulate binding of TNF- α to cell receptors, and/or modulate the consequential intracellular events comprising administering to a subject in a need thereof an effective amount of a benzobicyclooctane compounds of formula (I). The inhibition of TNF- α induced apoptosis and of NF κ B activation is one means of preventing and/or 25 treating autoimmune and inflammatory diseases including, but not limited to, rheumatoid arthritis, inflammatory bowel disease, psoriasis, atherosclerosis, asthma, reperfusion injury, ischemia, sepsis, graft vs. host disease, adult respiratory distress syndrome, multiple

sclerosis, and a host of severe invasive infections such as fulminant hepatitis, AIDS and bacterial meningitis, and allergic inflammation of the lungs and airways.

Thus, in one aspect, the present invention provides a method of inhibiting TNF- α induced apoptosis comprising administering to a subject in a need thereof an effective amount of a benzobicyclooctane compounds of formula (I). In another aspect, the present invention provides a method of inhibiting NF κ B activation comprising administering to a subject in a need thereof an effective amount of a benzobicyclooctane compounds of formula (I). In another aspect, the present invention provides a method of inhibiting, preventing, treating, or preventing and/or treating autoimmune and inflammatory diseases including, but not limited to, rheumatoid arthritis, Inflammatory Bowel Disease (IBD), psoriasis, atherosclerosis, asthma, reperfusion injury, ischemia, sepsis, graft vs. host disease, Adult Respiratory Distress Syndrome (ARDS), and multiple sclerosis, comprising administering to a subject in a need thereof an effective amount of a benzobicyclooctane compounds of formula (I). In another aspect, the present invention provides a method of inhibiting, preventing, treating, or preventing and/or treating severe invasive infections such as fulminant hepatitis comprising administering to a subject in a need thereof an effective amount of a benzobicyclooctane compounds of formula (I).

In another aspect, the present invention provides a method for the inhibition of IL-8 or other CXC chemokines binding to CXCR1 and/or CXCR2 receptors comprising administering an effective amount of a compound of formula (I) to a subject in need thereof. In another aspect, the present invention provides a method for reducing the levels of IL-8 within a subject comprising administering to a subject in need thereof an effective amount of a compound of formula (I). In another aspect, the present invention provides a method for treating, preventing, or treating and/or preventing one or more of inflammatory and autoimmune diseases such as Inflammatory Bowel Disease (IBD), psoriasis, rheumatoid arthritis, Acute Respiratory Distress Syndrome (ARDS), cancer, atherosclerosis, reperfusion injury, and graft vs. host disease, comprising administering to a subject in need thereof an effective amount of a compound of formula (I).

The present invention provides a method for inhibiting TNF- α mediated processes, comprising administering to a patient in need thereof, through a therapeutically or prophylactically acceptable manner, a therapeutically or pharmaceutically effective amount of a composition comprising a compound of formula (I). Administering may be by, for example, transdermal, oral, intravenous, intramuscular, vaginal, rectal, pulmonary, subcutaneous, sublingual and transmucosal administration.

The present invention provides a method for treating an inflammation event, comprising administering to a patient in need thereof, through a therapeutically or prophylactically acceptable manner, a therapeutically or pharmaceutically effective amount of the compound of formula (I). Administering may be selected from transdermal, oral, intravenous, intramuscular, vaginal, rectal, pulmonary, subcutaneous, sublingual and transmucosal administration.

The "effective amount" or "therapeutically effective amount" of a compound of the present invention will depend on the route of administration, the type of mammal being treated, and the physical characteristics of the specific mammal under consideration. These factors and their relationship to determining this amount are well known to skilled practitioners in the medical arts. This amount and the method of administration can be tailored to achieve optimal efficacy but will depend on such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize.

In addition, this invention provides a method for identifying a binding partner to a compound of formula (I), wherein the method comprises immobilizing proteins known to be involved in the TNF- α signaling pathway onto a suitable carrier; and passing a solution of said compounds in isolation or mixture over said proteins and analyzing for compound:protein complex formation using surface plasmon resonance (SPR) in a manner similar to that reported by Karlsson, R et al. Biosensor Analysis of Drug-Target Interactions: Direct and Competitive Binding Assays for Investigation of Interactions Between Thrombin and Thrombin Inhibitors. *Anal. Biochem.* **2000**, 278(1), 1-13. For other examples of identifying small molecule-protein interactions using SPR see the Biacore website: <http://www.biacore.com>

In addition, this invention provides a method for identifying a binding partner to a compound of formula (I), wherein the method comprises (in a manner similar to that reported by Shimizu, N et al. High Performance Affinity Beads for Identifying Drug Receptors. *Nature Biotechnology*, 2000, 18(8), 877-881) providing said compound(s)
5 bound to a solid support to provide solid phase compounds; contacting a cell or cell components with said solid phase compounds in isolation or mixture; removing uncomplexed cellular material, for example by gentle washing with aqueous buffer, from said solid phase compounds; and recovering said binding partner from the solid phase compounds.

10 As to each publication or patent referenced herein, that publication or patent is incorporated herein by reference in its entirety for all purposes.

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the
15 invention. Accordingly, the invention is not limited except as by the appended claims.


The following examples are offered by way of illustration, and not by way of limitation.

EXAMPLES

Abbreviations and acronyms used in the examples include: AcOH, acetic acid; APCI-MS, atmospheric pressure chemical ionization mass spectroscopy; DBU, 1,8-diazabicyclo[5.4.0]undec-7-ene; DEAD, diethylazodicarboxylate; DIEA, diisopropylethylamine; DMAP, 4-N,N-dimethylaminopyridine; DME, 1,2-dimethoxyethane; DMF, dimethylformamide; DMSO, dimethyl sulfoxide; DPPA, diphenylphosphoryl azide; ESI-MS, electrospray ionization mass spectroscopy; FAB-MS, fast atom bombardment mass spectroscopy; FTIR, Fourier transform infrared spectroscopy; HATU, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate; HPLC, high pressure liquid chromatography; HRMS, high resolution mass spectroscopy; LC-MS, liquid chromatography-mass spectroscopy; NMA, N-methylaniline; NMM, N-methylmorpholine; NMP, N-methylpyrrolidinone; NMR, nuclear magnetic resonance spectroscopy; TBAF, tetrabutylammonium fluoride; TFA, trifluoroacetic acid; THF, tetrahydrofuran; TSTU, O-(N-Succinimidyl)-1,1,3,3-tetramethyluronium tetrafluoroborate; rt, room temperature; h, hour; min, minute; eq, equivalents.

EXAMPLE 1

SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 9-(2-TRIMETHYLSILANYLETHYL) ESTER



A. 4-Hydroxy-11-oxo-tricyclo[6.2.2.0^{2,7}]dodeca-2(7),3,5-triene-9,10-dicarboxylic acid anhydride

A solution of dihydroxynaphthalene (500 g, 3.125 mol) and maleic anhydride (765 g, 7.815 mol, 2.5 eq) in 1 L 1:1 1,2-dichlorobenzene:toluene were heated at 110°C for 3 days. The reaction mixture was then cooled to 90°C, 1.5 L ethyl acetate added, and then further cooled to room temperature overnight. The mixture was then cooled over ice after another 0.5 L ethyl acetate was added and left stirring for 2 hours. The resultant solid was isolated by filtration, washed with 2x200 mL cold ethyl acetate and dried in oven at 40°C to provide 130 g of the anhydride as a beige solid (16% yield). ¹H NMR (acetone-*d*₆) 7.33 (d, *J*=8.2 Hz, 1H), 6.97 (d, *J*=2.3 Hz, 1H), 6.85 (dd, *J*=8.1, 2.4 Hz, 1H), 4.05 (d, *J*=3.9 Hz, 1H), 3.98 (dd, *J*=6.0, 2.9 Hz, 1H), 3.88 (dd, *J*=10.2, 3.9 Hz, 1H), 3.70 (dd, *J*=10.2, 2.7 Hz, 1H), 2.39 (br s, 2H). HRMS for MH⁺ 259.0600 (theoretical 259.0606).

B. 4-Hydroxy-11-oxo-tricyclo[6.2.2.0^{2,7}]dodeca-2(7),3,5-triene-9,10-dicarboxylic acid 9-(2-trimethylsilanylethyl) ester, 1


DMAP (0.96 g, 7.9 mmol, 10 mol%) and trimethylsilyl-ethanol (12.43 mL, 0.087 mol, 1.1 eq) were added to a stirred suspension of the anhydride from step A (20.5 g, 0.079 mol) in 200 mL acetonitrile and heated to reflux for 7 hours. By HPLC there was some starting material present and the two regioisomers of the opened anhydride were present in a 1:1 ratio. The reaction mixture was cooled and dicyclohexylamine (15.71 mL, 0.079 mol) was added dropwise. A precipitate formed instantaneously but was left overnight. The resulting white salt (40.73 g, 93%) was filtered, suspended in water, acidified with 2 M HCl and extracted with ethyl acetate. An emulsion formed, but was removed by filtration before the layers could be separated, and the organic layer was dried and evaporated *in vacuo* to give a mixture of the regioisomeric acid-esters as a beige foam (25.59 g, 93%). A hazy solution of the solids (25.59 g, 0.068 mol) in 150 mL isopropanol was treated with isopropylamine (5.79 mL, 0.068 mol) and left stirring overnight. The precipitate was isolated by filtration yielding a white solid (12.44 g, 42% yield) as a 86/17 mixture of diastereomers. This solid was slurried in 48 mL isopropanol for 1.5 hours giving

a second white solid (10.69 g) as a 93/7 mixture of diastereomers. This salt was cracked as described above to give a white foam (6.76 g) which was then triturated in 33 mL 20% diethyl ether/toluene at -20°C. The resulting white solid was collected by filtration and washed with 10 mL cold solvent. This afforded **1** as a white solid (4.59 g, 30% overall yield from the anhydride) of 98.2% purity by HPLC. ¹H NMR (acetonitrile-*d*₃) 7.12 (d, *J*=8.0 Hz, 1H), 6.76 (d, *J*=2.5 Hz, 1H), 6.72 (dd, *J*=8.0, 2.5 Hz, 1H), 4.20-4.06 (m, 2H), 3.65 (d, *J*=3 Hz, 1H), 3.61 (br s, 1H), 3.22 (br d, *J*=11.0 Hz, 1H), 2.93 (br s, 1H), 2.85 (dd, *J*=18.3, 2.3 Hz, 1H), 2.04 (br d, 1H), 0.99-0.93 (m, 2H), 0.03 (s, 9H). MS for MNa⁺ 399.4. Elemental analysis for C₁₉H₂₄O₆Si, Theoretical: C, 60.62; H, 6.43. Found: C, 60.61; H, 6.58.

10

EXAMPLE 2

SYNTHESIS OF (4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-(2-TRIMETHYLSILANYL-ETHYL) ESTER)




DMAP (0.5 g, 4 mmol, 10 mol%) and trimethylsilylethanol (6.6 mL, 5.45 g, 46 mmol) were added to a stirred suspension of the anhydride (10.83 g, 42 mmol) from Example 1.A in 400 mL acetonitrile and heated to reflux for 6 h. The volatiles were evaporated, and the resulting foam was chromatographed on silica gel (20% acetonitrile/dichloromethane with 2% AcOH). Appropriate fractions were combined and dichloromethane and toluene were used to remove residual AcOH. Repeated trituration of the less polar product with ethyl ether provided 4.9 g (31%) of the title compound. ¹H NMR (acetonitrile-*d*₃) 7.13 (d, *J*=8.0 Hz, 1H), 6.75 (d, *J*=2.2 Hz, 1H), 6.72 (dd, *J*=8.0, 2.5 Hz, 1H), 4.17-4.11 (m, 2H), 3.64-3.62 (m, 1H), 3.60 (d, *J*=3.0 Hz, 1H), 3.22 (dd, *J*=11.8, 3.0 Hz, 1H), 2.97 (dt, *J*=11.8, 2.2 Hz, 1H), 2.87 (dd, *J*=18.7, 2.2 Hz, 1H), 2.08 (ddd,

$J=18.4, 3.3, 2.5$ Hz, 1H), 1.00-0.95 (m, 2H), 0.04 (s, 9H). MS 399.4 (MNa^+). Elemental for $\text{C}_{19}\text{H}_{24}\text{O}_6\text{Si}$: Theoretical, C, 60.62; H, 6.43. Found: C, 60.58; H, 6.57. In addition, repeated trituration of the more polar product provided 5.0 g (32%) of acid **1**.

EXAMPLE 3


5 SYNTHESIS OF (9, 10 CIS)-10-ALLYLOXYCARBONYLAMINO-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



To a solution of acid **1** (196 mg, 0.52 mmol) in THF (25 mL) was added DPPA (230 μL , 1.05 mmol), triethylamine (150 μL , 1.08 mmol), and allyl alcohol (360 μL , 5.3 mmol). The mixture was heated to reflux and held for 15 h. Upon cooling, the mixture was concentrated *in vacuo*, and the residue chromatographed, initially with 30% ethyl acetate/hexane followed by a second chromatography using 15% ethyl acetate/dichloromethane to afford a total of 65.8 mg (30%) of the title compound. ESI-MS m/z 454 (MNa^+).

EXAMPLE 4

SYNTHESIS OF 5-HYDROXY-10-{METHYL-[(2,4,6-TRIMETHOXY-BENZYL-CARBAMOYL)-METHYL]-CARBAMOYL}-12-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5

A. Sarcosine-2,4,6-trimethoxybenzylamide

- To a solution of N-Fmoc-sarcosine (5 g, 16 mmol) in dichloromethane (160 mL) containing 3A molecular sieves was added NMM (6 mL, 5.52 g, 55 mmol), HATU (7.33 g, 19 mmol) and 2,4,6-trimethoxybenzylamine hydrochloride (4.5 g, 19.2 mmol).
- 10 The resulting reaction mixture was allowed to stir at rt overnight. The sieves were filtered, the volatiles evaporated and ethyl acetate was added. Acid wash (0.1 N HCl, 3x300 mL) followed by sodium bicarbonate (5% solution, 1x300 mL) provided a solid precipitate, which was collected, washed with ethyl acetate, collected and air dried. The organic layer was concentrated to dryness to give a residue which was triturated with ethyl acetate to
- 15 provide an additional amount of the Fmoc derivative of 101: amount recovered 7.5 g (95%). ¹H NMR (CDCl₃) 7.77-7.27 (m, 8H), 6.35 (br s, 1H), 6.05 (br s, 2H), 4.49-4.05 (m, 5H), 3.93 (s, 2H), 3.74 (s, 9H), 2.99 (s, 3H). FAB-MS m/z 513 (MNa⁺), 491 (MH⁺).

- The isolated N-Fmoc-sarcosine-2,4,6-trimethoxybenzylamide (6.5 g, 13 mmol) was suspended in 25% pyrrolidine/chloroform (100 mL) and allowed to stir at rt for
- 20 50 min. The volatiles were then evaporated to give a pale yellow solid. Column chromatography (10% methanol/dichloromethane) provided the desired product 101 upon trituration with ethyl ether, wt. 3.1 g (88%). ¹H NMR (CDCl₃) 6.13 (d, 2H), 4.48 (d, J=5.5


Hz, 2H), 3.82 (s, 9H), 3.24 (s, 2H), 2.40 (s, 3H). Elemental for $C_{13}H_{20}N_2O_4$: Theoretical, C, 58.19; H, 7.51; N, 10.44. Found: C, 58.09; H, 7.66; N, 10.18.

B. 5-Hydroxy-10-{methyl-[(2,4,6-trimethoxy-benzylcarbamoyl)-methyl]-carbamoyl}-12-oxo-tricyclo[6.2.2.0^{2,7}]dodeca-2(7),3,5-triene-9-(2-trimethylsilanyl-ethyl) ester

5 To a solution of the carboxylic acid from Example 2 (4.5 g, 11.8 mmol) in dichloromethane (25 mL) was added NMM (3.2 mL, 2.9 g, 29 mmol), HATU (5.3 g, 13.9 mmol), 3A molecular sieves and sarcosine-2,4,6-trimethoxybenzylamide (3.1 g). The resulting solution was allowed to stir at rt under nitrogen overnight. The volatiles were then evaporated, ethyl acetate (300 mL) was added and the organic layer was washed with
10 0.1 N HCl (2x150 mL), 5% $NaHCO_3$ solution (1x100 mL) and brine (1x100 mL). The organic layer was dried ($MgSO_4$), filtered and the volatiles were evaporated to give a yellow foam. Column chromatography (90% ethyl acetate/hexane) provided the desired product, wt. 3.5 g (49%). 1H NMR ($CDCl_3$) 7.05-6.70 (m, 3H), 6.48 (t, 1H), 6.12, 6.05 (2s, 2H), 4.56-4.40 (m, 2H), 4.20-2.71 (m, 21H), 2.20-2.07 (m, 1H), 1.00-0.90 (m, 2H), 0.02 (s,
15 9H). Elemental for $C_{32}H_{42}N_2O_9Si$: Theoretical, C, 60.16; H, 7.04; N, 4.25. Found: C, 60.29; H, 6.93; N, 4.18.

EXAMPLE 5

SYNTHESIS OF 2,4,6-TRIMETHOXY-BENZYL-CARBAMOYL-METHYL-CARBAMOYL-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIEN-4-YLOXYMETHYL)-BENZOIC ACID ALLYL ESTER



A. Allyl 4-Hydroxymethylbenzoate


To a solution of 4-hydroxymethylbenzoic acid (0.5 g, 3.3 mmol) in CHCl_3 (10 mL) was added allyl bromide (0.6 mL, 0.84 g, 6.9 mmol) and diisopropylethylamine (1.3 mL, 0.96 g, 7.5 mmol). The resulting reaction mixture was allowed to reflux under nitrogen for 2.5 h. Upon cooling to rt, dichloromethane (50 mL) was added and the organic layer was washed with 0.1 N HCl (3x30 mL), 5% NaHCO_3 solution (1x30 mL) and brine (1x30 mL). Upon drying (MgSO_4), filtration of the drying agent and concentration, the resulting oily residue was chromatographed on silica gel (30% ethyl acetate/hexane) to give 440 mg (70%) of a colorless oil. ^1H NMR (CDCl_3) 8.06 (d, $J=8.2$ Hz, 2H), 7.44 (d, $J=8.2$ Hz, 2H), 6.11-5.98 (m, 1H), 5.42 (dd, $J=17.2, 1.5$ Hz, 1H), 5.30 (dd, $J=10.4, 1.2$ Hz, 1H), 4.83 (dd, $J=5.6, 1.2$ Hz, 2H), 4.78 (s, 2H), 1.80 (br s, 1H). MS 192 (M^+).

B. 5-(4-Allyloxycarbonyl-benzyloxy)-10-{methyl-[(2,4,6-trimethoxybenzylcarbamoyl)-methyl]-carbamoyl}-11-oxo-tricyclo[6.2.2.0^{2,7}]dodeca-2(7),3,5-triene-9-carboxylic acid-(2-trimethylsilanyl-ethyl) ester

To a cooled solution (ice bath) of allyl 4-hydroxymethylbenzoate (0.62 g, 3.2 mmol) and the compound prepared in Example 4 (1.33 g, 2 mmol) in anhydrous THF (40 mL) was added PPh_3 (1.34 g, 5.1 mmol) and DEAD (0.8 mL, 0.88 g, 5.1 mmol). The resulting reaction mixture was allowed to warm to room temperature and then allowed to reflux under N_2 for 0.5 h. Column chromatography of the concentrated residue (90% ethyl acetate/hexane) provided the title compound as a white foamy material, wt. 0.98 g (58%). ^1H NMR (CDCl_3) 8.09 (d, $J=8.2$ Hz, 2H), 7.49 (d, $J=8.0$ Hz, 2H), 7.16-6.46 (m, 4H), 6.11-5.98 (m, 3H), 5.42 (d, $J=17.2$ Hz, 1H), 5.30 (d, $J=10.4$ Hz, 1H), 5.11 (s, 2H), 5.83 (d, $J=5.6$ Hz, 2H), 4.56-4.37 (m, 2H), 4.17-3.9 (m, 2H), 3.81-3.70 (m, 12H), 3.55-2.05 (m, 8H), 0.97-0.89 (m, 2H), 0.02 (s, 9H). Elemental for $\text{C}_{43}\text{H}_{52}\text{N}_2\text{O}_{11}\text{Si}$: Theoretical, C, 64.48; H, 6.54; N, 3.50. Found: C, 64.18; H, 6.67; N, 3.28.

EXAMPLE 6

SYNTHESIS OF 4-(10-DIPENTYL CARBAMOYL-9-{METHYL-[(2,4,6-TRIMETHOXY-BENZYL CARBAMOYL)-METHYL]-CARBAMOYL}-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIEN-4-YLOXYMETHYL)-BENZOIC ACID ALLYL ESTER



5

To a solution of the diester prepared in Example 5 (0.5 g, 0.62 mmol) in anhydrous THF (14 mL) was added a 1.0 M solution of TBAF (1.0 mL, 1 mmol). The reaction mixture was allowed to stir at rt for 1.75 h, after which ethyl acetate (200 mL) was added. The organic layer was then washed with 0.1 N HCl (2 x 50 mL), brine (2x50 mL), dried (MgSO₄), filtered and concentrated to give a colorless oil. The free acid was dissolved in dichloromethane (12 mL) and HATU (0.29 g, 0.76 mmol); NMM (0.17 mL, 0.16 g, 1.55 mmol) and dipentylamine (.15 mL, 0.12 g, 0.7 mmol) were added. The resulting reaction mixture was then allowed to stir at rt under N₂ for 3 days, after which dichloromethane (300 mL) was added. The organic layer was then washed with 0.1 N HCl (2x100 mL), 5% solution of NaHCO₃ (2x50 mL), water (2x50 mL), brine (1x50 mL), dried (MgSO₄), filtered and the volatiles were evaporated to give a colorless oil. Column chromatography (5% methanol/dichloromethane) provided the desired product, wt. 303 mg (58%). ¹H NMR (CDCl₃) 8.08 (d, *J*=8.2 Hz, 2H), 7.50 (d, *J*=8.3 Hz, 2H), 7.18-6.72 (m, 3H), 6.11-5.98 (m, 3H), 5.41 (dd, *J*=17.2, 1.3 Hz, 1H), 5.30 (apt t, *J*=6.0, 4.4 Hz, 1H), 5.11 (s, 2H), 4.83 (d, *J*=5.6 Hz, 2H), 4.53-4.35 (m, 2H), 4.02 (d, *J*=15.4 Hz, 1H), 3.85-2.80 (m, 23H), 2.16 (d, *J*=18.5 Hz, 1H), 1.60-1.10 (m, 12H), 0.94-0.86 (m, 6H). Elemental for C₄₈H₆₁N₃O₁₀.methanol: Theoretical, C, 67.49; H, 7.51; N, 4.82. Found: C, 67.68; H, 7.45; N, 4.63.


10

15

20

EXAMPLE 7

SYNTHESIS OF 4-(10-DIPENTYL-CARBAMOYL-9-{METHYL-[(2,4,6-TRIMETHOXY-BENZYL-CARBAMOYL)-METHYL]-CARBAMOYL}-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIEN-4-YLOXYMETHYL)-BENZOIC ACID



5


A solution of allyl ester prepared in Example 6 (0.28 g, 0.33 mmol), tetrakis(triphenylphosphine) palladium (0) (27 mg, 23 μ mol) and N-methylaniline (75 μ L, 74 mg, 0.69 mmol) in dichloromethane (3.5 mL) was allowed to stir at rt for 1 h. The reaction mixture was then diluted with dichloromethane (50 mL) and washed with 0.1 N HCl solution (2x20 mL) and brine (2x20 mL). Upon drying (MgSO_4), filtration of the drying agent and concentration, the resulting oily residue was chromatographed on silica gel (5% methanol/dichloromethane) to provide the desired product as a white solid, wt. 160 mg (60%). ^1H NMR (CDCl_3) 8.14, 7.92 (2 d, $J=8.0$, 7.8 Hz, 2H), 7.54 (d, $J=7.9$ Hz, 2H), 7.31-6.89 (m, 4H), 6.12 (s, 2H), 5.15, 5.06 (2 br s, 2H), 4.57-4.39 (m, 2H), 4.08 (d, $J=15.5$ Hz, 1H), 3.91-3.52 (m, 13H), 3.30-2.86 (m, 7H), 2.38 (s, 2H), 2.20 (d, $J=18.4$ Hz, 1H), 1.59-0.80 (m, 18H). MS (ESI +ve) 800 (MH^+), 822 (MNa^+). Elemental for $\text{C}_{45}\text{H}_{57}\text{N}_3\text{O}_{10}$.methanol: Theoretical, C, 66.41; H, 7.39; N, 5.05. Found: C, 66.28; H, 7.22; N, 4.83.

10

15

EXAMPLE 8

SYNTHESIS OF 4-[4-(2-DIMETHYLCARBAMOYL-PYRROLIDINE-1-CARBONYL)-BENZYLOXY]-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-DIPENTYLAMIDE 10-{METHYL-[(2,4,6-TRIMETHOXY-BENZYL-CARBAMOYL)-METHYL]-AMIDE}



5


A solution of the acid prepared in Example 7 (67 mg, 84 μ mol), HATU (38 mg, 100 μ mol), NMM (22 μ L, 20 mg, 0.2 mmol) and H-prolinedimethylamide (15 mg, 0.1 mmol) in dichloromethane (1.0 mL) was allowed to stir at rt overnight. The reaction mixture was then diluted with dichloromethane (30 mL) and washed with 0.1 N HCl solution (2x20 mL), 5% NaHCO₃ solution (1x25 mL) and brine (1x25 mL). Upon drying (MgSO₄), filtration of the drying agent and concentration, the resulting oily residue was chromatographed on silica gel (5% methanol/dichloromethane) to provide the diastereomeric mixture as a white solid, wt. 54 mg (70%). ¹H NMR (CDCl₃) 7.61 (d, *J*=8.0 Hz), 7.43 (d, *J*=8.0 Hz), 7.38 (s), 7.15(d, *J*=8.0 Hz), 7.02-6.78 (m), 6.11 (s), 6.09 (s), 5.06 (s), 5.02 (s), 4.52-4.33 (m), 4.04-3.88 (m), 3.86-3.67 (m), 3.57-3.49 (m), 3.27-3.06 (m), 2.99 (s), 2.85 (s), 2.82 (s), 2.76 (d, *J*=10.6 Hz), 2.53 (d, *J*=9.1 Hz), 2.31-1.80 (m), 1.60-1.10 (m), 0.91 (t, *J*=5.8 Hz), 0.87 (t, *J*=6.0 Hz). ESI-MS *m/z* 925 (MH⁺), 947 (MNa⁺).

10

15

EXAMPLE 9

SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-(2,4-DIMETHOXY-BENZYL) ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5 To a solution of acid **1** (2.83 g, 7.5 mmol), prepared in Example 1, 2,4-dimethoxybenzylalcohol (1.65 g, 9.8 mmol) and DMAP (0.1 g, 0.8 mmol) in dichloromethane (50 mL) was added DIEA (2.8 mL, 2.1 g, 16 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (2.2 g, 11.5 mmol). The

10 resulting reaction mixture was allowed to stir at rt for 24 h, after which time it was concentrated to dryness, redissolved in ethyl acetate (300 mL) and washed with 0.1 N HCl solution (2x100 mL), 5% NaHCO₃ solution (1x100 mL) and brine (1x100 mL). Upon drying (MgSO₄), filtration of the drying agent and concentration, the resulting oily residue

15 was chromatographed on silica gel (35% ethyl acetate/hexane) to provide 370 mg (10%) of the trans bis-ester. ¹H NMR (CDCl₃) 7.10 (dd, *J*=8.1, 2.9 Hz, 2H), 6.69 (dd, *J*=8.0, 2.4 Hz, 1H), 6.48-6.42 (m, 2H), 6.33 (d, *J*=2.4 Hz, 1H), 5.30-5.22 (br s, 1H), 5.03 (dd, *J*=37.1, 11.8 Hz, 2H), 4.27-4.21 (m, 2H), 3.91 (d, *J*=2.2 Hz, 1H), 3.83 (s, 3H), 3.80 (s, 3H), 3.76-3.73 (m, 1H), 3.69 (dd, *J*=5.8, 2.2 Hz, 1H), 3.28-3.22 (m, 1H), 2.40 (dd, *J*=18.9, 2.0 Hz, 1H), 2.10 (dm, 1H), 1.04-0.99 (m, 2H), 0.01 (s, 9H). FAB-MS *m/z* 526 (M⁺).

EXAMPLE 10

SYNTHESIS OF 4-METHOXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-(2,4-DIMETHOXY-BENZYL) ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5


To a solution of the phenol prepared in Example 9 (70 mg, 0.13 mmol) in anhydrous THF (5 mL) was added cesium carbonate (48 mg, 0.14 mmol) and methyl iodide (45 μ L, 103 mg, 0.7 mmol). The resulting reaction mixture was allowed to stir at rt under N₂ for 22 h. The reaction mixture was then diluted with ethyl acetate (20 mL) and washed with 0.1 N HCl solution (2x10 mL), 5% NaHCO₃ solution (1x10 mL) and brine (1x10 mL). Upon drying (MgSO₄), filtration of the drying agent and concentration, the resulting oily residue was chromatographed on silica gel (1% methanol/dichloromethane) to provide 35 mg (49%) of the methyl ether. ¹H NMR (CDCl₃) 7.18 (d, *J*=8.2 Hz, 1H), 7.11 (d, *J*=8.2 Hz, 1H), 6.77 (dd, *J*=8.2, 2.5 Hz, 1H), 6.58 (d, *J*=2.4 Hz, 1H), 6.49-6.41 (m, 2H), 5.02 (dd, *J*=24.7, 11.8 Hz, 2H), 4.28-4.22 (m, 2H), 3.98 (d, *J*=2.2 Hz, 1H), 3.83 (s, 3H), 3.80 (s, 3H), 3.78-3.75 (m, 1H), 3.72 (s, 3H), 3.71-3.69 (m, 1H), 3.28-3.26 (m, 1H), 2.41 (dd, *J*=19.0, 2.1 Hz, 1H), 2.11 (dm, 1H), 1.05-0.99 (m, 2H), 0.06 (s, 9H). FAB-MS *m/z* 540 (M⁺).

10

15

EXAMPLE 11

SYNTHESIS OF 10-(2,4-DIMETHOXY-BENZYL-CARBAMOYL)-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5 General Procedure A for the Synthesis of 4-Hydroxy-10-amido Derivatives


To a solution of the cis acid ester **1** (0.5 mmol) and molecular sieves (3A) in THF (2 mL) was added diisopropylethylamine (2.8 mmol) and diphenylphosphoryl azide (0.7 mmol). The solution was allowed to stir at rt under nitrogen for 3-4 h, after which time a selected amine (1.5 – 2 mol equivalents) and DMAP (2 mol equivalents) were added and the resulting reaction mixture was allowed to stir overnight. Dilution with ethyl acetate (25 mL), followed by washes with 1 N HCl (2x25 mL), 5% NaHCO₃ solution (2x25 mL) and brine (1x25 mL) provided a pale yellow solution, which was dried (MgSO₄), filtered and concentrated to dryness. Column chromatography provided the desired product.

The title compound was prepared as in general procedure A, above. Column chromatography (10% acetonitrile/dichloromethane) provided 35% of the title compound. ¹H NMR (acetonitrile-*d*₃) 7.14 (br s, 1H), 7.11 (d, *J*=8.2 Hz, 1H), 7.05 (br s, 1H), 7.02 (d, *J*=8.2 Hz, 1H), 6.65 (dd, *J*=8.0, 2.5 Hz, 1H), 6.53 (dd, *J*=8.4, 2.3 Hz, 2H), 6.44 (dd, *J*=8.4, 2.3 Hz, 1H), 4.25-4.16 (m, 4H), 3.81, 3.77 (2 s, 6H), 3.74 (app. q, *J*=2.4 Hz, 1H), 3.69 (d, *J*=1.9 Hz, 1H), 3.39 (dd, *J*=6.3, 1.9 Hz, 1H), 3.20 (dt, *J*=6.3, 2.2 Hz, 1H), 2.37 (dd, *J*=19.0, 2.2 Hz, 1H), 2.03 (dq, *J*=19.0, 3.0, 2.2 Hz, 1H), 1.01-0.96 (m, 2H), 0.04 (s, 9H). ¹³C NMR (acetonitrile-*d*₃) 209.20, 174.55, 171.38, 161.73, 159.66, 157.60, 135.53, 134.32, 130.69, 126.08, 120.02, 115.50, 114.98, 105.26, 99.43, 64.48, 57.56, 56.30, 56.10, 46.27,

43.91, 39.42, 39.01, 38.41, 17.96, -1.43. FAB-MS m/z 540 (MH^+). Elemental for $C_{28}H_{35}NO_7Si$: Theoretical, C, 63.98; H, 6.71; N, 2.66. Found: C, 63.77; H, 6.86; N, 2.63.

EXAMPLE 12


5 SYNTHESIS OF 10-DIPENTYLCARBAMOYL-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as in general procedure A in Example 11. Column chromatography (5% acetonitrile/dichloromethane) provided 23% of the title compound. 1H NMR (acetonitrile- d_3) 7.12 (d, $J=8.0$ Hz, 1H), 6.98 (br s, 1H), 6.67 (dd, $J=8.0, 2.5$ Hz, 1H), 6.53 (d, $J=2.5$ Hz, 1H), 4.62-4.23 (m, 2H), 3.76-3.73 (m, 1H), 3.66 (dd, $J=6.6, 1.7$ Hz, 1H), 3.56-3.19 (m, 5H), 2.97-2.88 (m, 1H), 2.47 (dd, $J=18.8, 2.1$ Hz, 1H), 2.04 (dt, $J=19.0, 2.6$ Hz, 1H), 1.69-1.15 (m, 12H), 1.02-0.97 (m, 2H), 0.92, 0.88 (2 t, $J=5.9, 5.6$ Hz, 6H), 0.03 (s, 9H). ^{13}C NMR (acetonitrile- d_3) 208.83, 174.55, 170.74, 157.60, 135.21, 134.56, 125.88, 115.47, 114.94, 64.27, 57.27, 48.65, 46.87, 46.77, 40.21, 38.98, 38.25, 29.94, 29.86, 29.75, 28.34, 23.30, 23.26, 18.10, 14.44, -1.46. FAB-MS m/z 516 (MH^+). Elemental for $C_{29}H_{45}NO_5Si$: Theoretical, C, 67.53; H, 8.79; N, 2.72. Found: C, 67.36; H, 9.00; N, 2.73.

EXAMPLE 13

SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5 A mixture of the product from Example 1 (1.46g, 3.88 mmol), TSTU (1.40g, 4.65 mmol), DIEA (3.2 mL, 18.4 mmol) were dissolved in THF (15 mL) and stirred under N₂ for 5 h. The solution was then treated with DMAP (0.58g, 4.7 mmol) and n-propanol (7.5 mL) and stirred an additional 19 h. The reaction was quenched with 0.2 M HCl (aq) and diluted with 150 mL ethyl acetate. The phases were separated and the organic was

10 washed with 5% NaHCO₃ (aq) and brine. The organic layer was separated, dried (Na₂SO₄) and concentrated to 2.0g of light yellow oil. Silica chromatography (ethyl acetate/hexanes) afforded 1.0 g (62%) of the title compound. ¹H NMR (CDCl₃) 7.12 (d, *J*=8 Hz, 1H), 6.70 (dd, *J*=2.5, 8 Hz, 1H), 6.64 (d, *J*=2.5 Hz, 1H), 5.26 (s, 1H), 4.25 (dd, *J*=7, 9 Hz, 2H), 3.98 – 3.93 (m, 3H), 3.74 (d, *J*=2.5 Hz, 1H), 3.65 (dd, *J*=2.2, 5.8 Hz, 1H), 3.2 (dd, *J*=2, 5 Hz,

15 1H), 2.41 (dd, *J*=2, 19 Hz, 1H), 2.16 (dd, *J*=2, 21 Hz, 1H), 1.57 (dd, *J*=7, 14 Hz, 2H), 1.05 (ddd, *J*=7, 7, <1 Hz, 2H), 0.87 (t, *J*=7, 7 Hz, 3H), 0.04 (s, 9H); ESI-MS *m/z* 417 (M-H)⁺.

EXAMPLE 14

SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-(2-CYCLOHEXYLOXY-ETHYL) ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



General Procedure B for the Synthesis of Trans Bis-esters 3


To a solution of the cis acid ester **1** (0.27 mmol) and molecular sieves (3A) in THF (1.0 mL) was added DIEA (1.4 mmol) and DPPA (0.37 mmol). The solution was allowed to stir at rt under nitrogen for 3-4 h, after which time a selected alcohol (2.5 mol equivalents) and DMAP (2 mol equivalents) were added and the resulting reaction mixture was allowed to stir overnight. Dilution with ethyl acetate (25 mL), followed by washes with 1 N HCl (2x25 mL), 5% NaHCO₃ solution (2x25 mL) and brine (1x25 mL) provided a pale yellow solution, which was dried (MgSO₄), filtered and concentrated to dryness. Column chromatography provided the desired product.

The title compound was prepared as described in general procedure B using 2-cyclohexyloxyethanol. Column chromatography (7% acetonitrile/dichloromethane) provided a 30% yield of the title compound. ¹H NMR (acetonitrile-*d*₃) 7.15 (d, *J*=8.0 Hz, 1H), 7.01 (br s, 1H), 6.74-6.69 (m, 2H), 4.32-4.19 (m, 2H), 4.14-4.05 (m, 2H), 3.84 (d, *J*=2.2 Hz, 1H), 3.73 (app. q, *J*=2.7 Hz, 1H), 3.60 (dd, *J*=6.0, 2.2 Hz, 1H), 3.56-3.53 (m, 2H), 3.27-3.20 (m, 1H), 3.09 (dt, *J*=6.0, 2.3 Hz, 1H), 2.36 (dd, *J*=19.0, 2.5 Hz, 1H), 2.06 (ddd, *J*=19.0, 3.2, 2.1 Hz, 1H), 1.87-1.75 (m, 2H), 1.74-1.64 (m, 2H), 1.56-1.46 (m, 1H), 1.34-1.12 (m, 5H), 1.06-1.00 (m, 2H), 0.05 (s, 9H). ¹³C NMR (acetonitrile-*d*₃) 208.86, 173.97, 172.62, 157.71, 135.64, 134.36, 126.38, 115.77, 115.03, 78.49, 66.37, 65.95, 64.74,

55.95, 47.07, 43.78, 39.32, 38.30, 33.03, 26.63, 24.84, 18.01, -1.42. FAB-MS m/z 502 (M^+). Elemental for $C_{27}H_{38}O_7Si$: Theoretical, C, 64.51; H, 7.62. Found: C, 64.47; H, 7.76.

EXAMPLE 15


SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-
5 DICARBOXYLIC ACID 10-(2-PYRIDIN-2-YL-ETHYL) ESTER 9-(2-TRIMETHYLSILANYL-ETHYL)
ESTER



The title compound was prepared as described in general procedure B using 2-(2-hydroxyethyl)-pyridine. Column chromatography (neat ethyl acetate) provided 48% of
10 the title compound. 1H NMR (acetonitrile- d_3) 8.49 (ddd, $J=4.7, 1.7, 1.1$ Hz, 1H), 7.68 (td, $J=7.7, 1.9$ Hz, 1H), 7.23-7.18 (m, 2H), 7.14 (d, $J=8.2$ Hz, 1H), 6.72 (dd, $J=8.0, 2.5$ Hz, 1H), 6.57 (d, $J=2.5$ Hz, 1H), 4.42-4.15 (2 m, 4H), 3.72-3.68 (m, 2H), 3.53 (dd, $J=6.0, 2.2$ Hz, 1H), 3.04-2.99 (m, 3H), 2.33 (dd, $J=19.0, 2.2$ Hz, 1H), 2.03 (ddd, $J=19.0, 3.2, 2.1$ Hz, 1H), 1.03-0.97 (m, 2H), 0.04 (s, 9H). ^{13}C NMR (acetonitrile- d_3) 208.85, 173.90, 172.50, 159.30, 157.78,
15 150.50, 137.92, 135.53, 134.23, 126.38, 124.73, 123.06, 115.82, 114.95, 65.40, 64.68, 55.86, 46.87, 43.75, 39.22, 38.30, 37.71, 17.95, -1.42. FAB-MS m/z 482 (MH^+). Elemental for $C_{26}H_{31}NO_6Si$: Theoretical, C, 64.84; H, 6.49; N, 2.91. Found: C, 64.63; H, 6.43; N, 2.70.

EXAMPLE 16

SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-(3-FLUORO-BENZYL) ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5 The title compound was prepared as described in general procedure B using 3-fluorobenzyl alcohol. Column chromatography (1% methanol/dichloromethane) provided a 56% yield of the title compound. ¹H NMR (acetonitrile-*d*₃) 7.41-7.34 (m, 1H), 7.16 (d, *J*=8.2 Hz, 1H), 7.11-7.02 (m, 3H), 6.98 (br s, 1H), 6.72 (dd, *J*=8.2, 2.5 Hz, 1H), 6.61 (d, *J*=2.5 Hz, 1H), 5.03 (s, 2H), 4.27-4.22 (m, 2H), 3.86 (d, *J*=2.2 Hz, 1H), 3.74 (dd, *J*=5.2, 2.5 Hz, 1H), 3.67 (dd, *J*=6.0, 2.2 Hz, 1H), 3.12 (dt, *J*=6.0, 2.3 Hz, 1H), 2.38 (dd, *J*=19.0, 2.2 Hz, 1H), 2.06 (ddd, *J*=19.1, 3.3, 2.2 Hz, 1H), 1.03-0.97 (m, 2H), 0.04 (s, 9H). ¹³C NMR (acetonitrile-*d*₃) 208.68, 173.91, 172.49, 165.58, 162.35, 157.72, 140.06, 139.95, 135.46, 134.36, 131.72, 131.62, 126.43, 124.97, 124.94, 116.26, 115.99, 115.91, 115.82, 115.61, 114.97, 67.07, 64.74, 55.81, 47.06, 43.81, 39.19, 38.27, 17.96, -1.45. FAB-MS *m/z* 484 (M⁺). Elemental for C₂₆H₂₉FO₆Si: Theoretical, C, 64.44; H, 6.03. Found: C, 64.47; H, 6.13.

10

15

EXAMPLE 17

SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-(2-PYRROLIDIN-1-YL-ETHYL) ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5


The title compound was prepared as described in general procedure B using 1-(2-hydroxyethyl)-pyrrolidine. Column chromatography (10% methanol/dichloromethane) provided 39% of the title compound. ¹H NMR (acetonitrile-*d*₃) 7.17-7.13 (m, 1H), 6.73-6.68 (m, 2H), 4.29-4.21 (m, 2H), 4.08 (t, *J*=5.8 Hz, 2H), 3.83 (d, *J*=2.2 Hz, 1H), 3.72 (dd, *J*=5.5, 2.8 Hz, 1H), 3.57 (dd, *J*=6.0, 2.2 Hz, 1H), 3.11 (dt, *J*=6.0, 2.3 Hz, 1H), 2.63 (td, *J*=5.6, 2.5 Hz, 2H), 2.51-2.45 (m, 4H), 2.35 (dd, *J*=19.0, 2.2 Hz, 1H), 2.05 (ddd, *J*=19.0, 3.2, 2.1 Hz, 1H), 1.73-1.68 (m, 4H), 1.05-1.00 (m, 2H), 0.05 (s, 9H). ¹³C NMR (acetonitrile-*d*₃) 208.89, 174.00, 172.59, 157.89, 135.58, 134.18, 126.41, 126.34, 115.83, 115.09, 114.94, 65.13, 64.69, 55.92, 55.10, 55.04, 54.93, 46.99, 43.79, 43.72, 39.27, 38.33, 24.31, 24.11, 17.99, -1.42. FAB-MS *m/z* 474 (MH⁺).

10

15

EXAMPLE 18


SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-DODECYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 The title compound was prepared as described in general procedure B using n-dodecanol. Column chromatography (1% methanol/dichloromethane) provided 50% of the title compound. ¹H NMR (acetonitrile-*d*₃) 7.15 (d, *J*=8.0 Hz, 1H), 7.00 (br s, 1H), 6.71 (dd, *J*=8.0, 2.5 Hz, 1H), 6.67 (d, *J*=2.5 Hz, 1H), 4.28-4.22 (m, 2H), 4.04-3.91 (m, 2H), 3.83 (d, *J*=2.2 Hz, 1H), 3.72 (dd, *J*=5.4, 2.6 Hz, 1H), 3.57 (dd, *J*=5.9, 2.3 Hz, 1H), 3.09 (dt, *J*=6.0, 2.4 Hz, 1H), 2.37 (dd, *J*=19.0, 2.5 Hz, 1H), 2.05 (ddd, *J*=19.0, 3.1, 2.1 Hz, 1H), 1.56-1.44 (m, 2H), 1.28 (br s, 18H), 1.05-1.00 (m, 2H), 0.88 (t, *J*=6.6 Hz, 3H), 0.05 (s, 9H). ¹³C NMR (acetonitrile-*d*₃) 208.88, 174.01, 172.64, 157.75, 135.65, 134.36, 126.39, 115.75, 114.92, 66.38, 64.68, 55.98, 47.05, 43.87, 39.27, 38.32, 32.78, 30.52, 30.49, 30.44, 30.35, 30.21, 30.03, 29.41, 26.69, 23.50, 18.01, 14.50, -1.40. ESI-MS *m/z* 567.3 (MNa⁺).
- 10
- 15 Elemental for C₃₁H₄₈O₆Si: Theoretical, C, 68.34; H, 8.88. Found: C, 68.22; H, 8.98.

EXAMPLE 19

SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-ALLYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5 The title compound was prepared as described in general procedure B using allyl alcohol. Column chromatography (7% acetonitrile/dichloromethane) provided 39% of the title compound. ¹H NMR (acetonitrile-*d*₃) 7.16 (d, *J*=8.0 Hz, 1H), 7.02 (br s, 1H), 6.72 (dd, *J*=8.0, 2.5 Hz, 1H), 6.69 (d, *J*=2.5 Hz, 1H), 5.92-5.81 (m, 1H), 5.28-5.17 (m, 2H), 4.56-4.43 (m, 2H), 4.33-4.19 (m, 2H), 3.87 (d, *J*=2.2 Hz, 1H), 3.74 (dd, *J*=5.4, 2.6 Hz, 1H), 3.64 (dd, *J*=5.9, 2.3 Hz, 1H), 3.12 (dt, *J*=5.9, 2.5 Hz, 1H), 2.37 (dd, *J*=19.0, 2.2 Hz, 1H), 2.06 (ddd, *J*=19.0, 3.3, 2.2 Hz, 1H), 1.06-1.00 (m, 2H), 0.05 (s, 9H). ¹³C NMR (acetonitrile-*d*₃) 208.79, 173.96, 172.37, 157.71, 135.58, 134.36, 133.51, 126.44, 118.75, 115.79, 114.95, 66.75, 64.72, 55.92, 47.01, 43.81, 39.22, 38.31, 17.98, -1.43. FAB-MS *m/z* 416 (*M*⁺). Elemental for C₂₂H₂₈O₆Si: Theoretical, C, 63.44; H, 6.78. Found: C, 63.19; H, 6.97.

15

EXAMPLE 20


SYNTHESIS OF 10-AZIDOCARBONYL-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



To a solution of 1 (584 mg, 1.51 mmol) in THF (15.6 mL) was added triethylamine (0.864 mL, 6.14 mmol) and DPPA (0.384 mL, 1.79 mmol). After 6.5 h the reaction was diluted with ethyl acetate and 1% HCl. The layers were separated and the organic layer washed with 5% NaHCO₃, H₂O, and brine, then dried (Na₂SO₄) and concentrated. Flash chromatography (25% ethyl acetate/hexane) afforded the trans acyl azide 2 (218 mg, 36%). FTIR (NaCl, cm⁻¹): 3435, 2960, 2907, 2152, 2129, 1803, 1735, 1728. ¹H NMR (CDCl₃): 7.15 (d, 1H), 6.7 (m, 2H), 4.23 (m, 2H), 3.94 (d, 1H), 3.77 (dd, 1H), 3.68 (dd, 1H), 3.20 (ddd, 1H), 2.40 (dd, 1H), 2.15 (ddd, 1H), 0.10 (s, 9H).

EXAMPLE 21

10 SYNTHESIS OF 4-HYDROXY-11-OXO-10-PROPOXYCARBONYLAMINO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




General Procedure C for Synthesis of 10-Alkoxycarbonylamino Derivatives.

A 0.1 M solution of the acyl azide 2 in dioxane was refluxed for 1 h to generate the isocyanate. A selected alcohol was added in large excess and the solution heated for 4-20 h. The solution was cooled, and the crude product isolated by concentration *in vacuo* or by extraction. Chromatography on SiO₂ afforded the product.

The title compound was prepared as described in general procedure C employing propyl alcohol. Chromatography with 10% to 30% ethyl acetate/hexane afforded the title compound in 11% overall yield. FAB-MS m/z 434 (MH⁺).

EXAMPLE 22

SYNTHESIS OF 4-HYDROXY-10-(5-METHYL-ISOXAZOL-3-YLMETHOXYCARBONYLAMINO)-11-
OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5 General Procedure D for Synthesis of 10-Alkoxycarbonylamino Derivatives.

A 0.1 M solution of the acyl azide **2** in dioxane was refluxed for 1 h to generate the isocyanate. The alcohol (1.2 - 2 eq) was added followed by DMAP and the solution refluxed until complete, typically 12-20 h. The crude product was isolated by extraction.

The title compound was prepared as described in general procedure D employing 5-methylisoxazole-3-methanol, and with the following modifications. A solution of the acyl azide (200 mg, ~0.36 mol) in dioxane (3.6 mL) was heated to reflux for 30 min. The reaction was cooled to rt and 171 mg of 5-methylisoxazole-3-methanol was added. The reaction was returned to reflux for 15 h, then cooled to rt. The reaction was quenched with aqueous ammonium chloride and diluted with ethyl acetate. The phases were partitioned, and the organic layer was separated and washed with 5% aqueous NaHCO₃, then brine. The solution was dried (Na₂SO₄) and concentrated to dryness. Chromatography with 5% to 40% ethyl acetate/hexane afforded the title compound in 4% overall yield. ¹H NMR (CDCl₃) 7.12 (d, 1H), 6.71 (dd, 1H), 6.56 (d, 1H), 5.81 (s, 1H), 5.49 (s, 1H), 4.97 (dd, 2H), 4.25 (dd, 2H), 3.95 (d, 1H), 3.75 (m, 2H), 3.48 (s, 1H), 3.25 – 3.23 (m, 1H), 2.41 (s, 3H), 2.38 (dd, 1H), 2.25 (dd, 1H), 1.03 (dd, 2H), 0.05 (s, 9H).

EXAMPLE 23


SYNTHESIS OF 4-HYDROXY-10-ISOPROPOXYCARBONYLAMINO-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5 The title compound was prepared as described in general procedure C employing isopropyl alcohol. Chromatography with 30% ethyl acetate/hexane afforded a 53% yield of product. ESI-MS m/z 456 (MNa^+).

EXAMPLE 24


10 SYNTHESIS OF 10-CYCLOPENTYLOXYCARBONYLAMINO-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as in general procedure C employing cyclopentanol. Chromatography with 30% ethyl acetate/hexane afforded a 64% yield of product. ESI-MS m/z 482 (MNa^+).

EXAMPLE 25


SYNTHESIS OF (9, 10 TRANS)-10-ALLYLOXYCARBONYLAMINO-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 The title compound was prepared as in general procedure C employing allyl alcohol. Chromatography with 40% ethyl acetate/hexane afforded a 44% yield of product. ESI-MS m/z 454 (MNa⁺).

EXAMPLE 26


10 SYNTHESIS OF 4-HYDROXY-10-(INDAN-2-YLOXYCARBONYLAMINO)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as in general procedure D employing 2-indanol. Chromatography with 30% ethyl acetate/hexane afforded 17% yield of product. ESI-MS m/z 530 (MNa⁺).

EXAMPLE 27

SYNTHESIS OF 10-(3-ALLYL-UREIDO)-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5 General Procedure E for The Synthesis of 10-Ureido Derivatives

A 0.1 M solution of the acyl azide **2** in dioxane was refluxed for 0.5 h. Upon cooling the appropriate amine was added and the solution stirred at ambient temperature for 1-4 h. Ethyl acetate and 1% HCl were added and the layers separated. The organic layer was washed with 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄) and concentrated *in vacuo*. Flash chromatography afforded the urea.

The title compound was prepared as described in general procedure E employing 3.6 eq of allylamine. The crude product was chromatographed on SiO₂ to afford a 27% yield of the urea. ESI-MS *m/z* 453 (MNa⁺).

EXAMPLE 28

15 SYNTHESIS OF 4-HYDROXY-10-{3-[2-(4-HYDROXY-PHENYL)-ETHYL]-UREIDO}-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




The title compound was prepared as described in general procedure E employing 2.2 eq of tyramine. The crude product was chromatographed with 50% ethyl

acetate/hexane to 65% ethyl acetate/hexane gradient to afford an 18% yield of the urea. ESI-MS m/z 533 (MNa^+).

EXAMPLE 29

5 SYNTHESIS OF 4-HYDROXY-10-[(MORPHOLINE-4-CARBONYL)-AMINO]-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




The title compound was prepared as described in general procedure E employing 3.3 eq of morpholine. The crude product was chromatographed with 70% ethyl acetate/methylene chloride to afford a 50% yield of the urea. ESI-MS m/z 483 (MNa^+).

10

EXAMPLE 30


SYNTHESIS OF 10-(3-TERT-BUTYL-UREIDO)-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



15 The title compound was prepared as described in general procedure E employing 3 eq of tert-butylamine. The crude product was chromatographed with 35% ethyl acetate/hexane to 45% ethyl acetate/hexane gradient to afford a 57% yield of the urea. ESI-MS m/z 469 (MNa^+).

EXAMPLE 31


SYNTHESIS OF 10-[3-(2,4-DIMETHOXY-BENZYL)-UREIDO]-4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5 The title compound was prepared as described in general procedure E employing 1.5 eq of 2,4-dimethoxybenzylamine. The crude product was chromatographed with 60% ethyl acetate/hexane to afford a 16% yield of the urea. ESI-MS m/z 563 (MNa^+).

EXAMPLE 32


10 SYNTHESIS OF 4-HYDROXY-10-(3-NAPHTHALEN-1-YLMETHYL-UREIDO)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as described in general procedure E employing 2 eq of 1-naphthalene methylamine. The crude product was chromatographed with 50% ethyl acetate/hexane to afford a 47% yield of the urea. ESI-MS m/z 531 (MH^+).

EXAMPLE 33

SYNTHESIS OF 4-(4-ALLYLOXYCARBONYL-BENZYLOXY)-11-OXO-
TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER
9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5


General Procedure F for the Synthesis of 4-Alkoxy Derivatives

To a 0.1 M solution of the benzobicyclooctane phenol (1 eq), prepared in Example 13, in tetrahydrofuran was added triphenylphosphine and the appropriate alcohol. The solution was cooled to 0°C and DEAD was added. The cooling bath was removed, the solution stirred at ambient temp for 5 min then heated at reflux until reaction was complete, typically 20-30 min. After cooling, the solution was diluted with ethyl acetate, water was added and the layers separated. The organic layer was washed with brine, dried (Na₂SO₄), and concentrated *in vacuo*. Chromatography on SiO₂ afforded the aryl ether.

The title compound was prepared as described in general procedure F employing 2.5 eq triphenylphosphine, 1.9 eq of allyl-4-(hydroxymethyl)-benzoate, and 2.5 eq of DEAD. Chromatography with 15% ethyl acetate/hexane followed by a second chromatography with 25% ethyl acetate/hexane afforded a 54% yield of aryl ether. ESI-MS m/z 615 (MNa⁺).

EXAMPLE 34

SYNTHESIS OF 4-[4-(2-DIMETHYLCARBAMOYL-PYRROLIDINE-1-CARBONYL)-BENZYLOXY]-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5

A. 4-(4-Carboxy-benzyloxy)-11-oxo-tricyclo[6.2.2.0^{2,7}]dodeca-2(7),3,5-triene-9,10-dicarboxylic acid 10-propyl ester 9-(2-trimethylsilyl-ethyl) ester

To a solution of the allyl ester prepared in Example 33 in methylene chloride was added N-methylaniline (40 μ L, 0.37 mmol) followed by tetrakis(triphenylphosphine)palladium(0) (21 mg, 0.02 mmol). The reaction was stirred for 20 min, diluted with ethyl acetate, and 2% HCl added. The layers were separated and the organic layer was washed with 1% HCl, H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*. Chromatography with 65% ethyl acetate/methylene chloride to 95% ethyl acetate/dichloromethane afforded a 60% yield of product. ESI-MS m/z 551 (M-H)⁻.

15 B. 4-[4-(2-Dimethylcarbamoyl-pyrrolidine-1-carbonyl)-benzyloxy]-11-oxo-tricyclo[6.2.2.0^{2,7}]dodeca-2(7),3,5-triene-9,10-dicarboxylic acid 10-propyl ester 9-(2-trimethylsilyl-ethyl) ester


To a solution of the acid (51.1 mg), N-methylmorpholine (30 μ L), and proline dimethylamide (17.5 mg, 0.12 mmol) in methylene chloride (0.7 mL) was added HATU (46 mg, 0.12 mmol). The solution was stirred for 6 h, diluted with ethyl acetate, and quenched with 3% HCl. The layers were separated and the organic layer washed with 1% HCl, 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*.

Chromatography with 100% ethyl acetate - 5% methanol/methylene chloride gradient afforded the product contaminated with tetramethyl urea. An ethereal solution of the mixture was washed with H₂O (10X), dried (Na₂SO₄) and concentrated to afford 29.6 mg (46%) of the titled product as a 1/1 mixture of diastereomers. ESI-MS m/z 677 (MH⁺).

5

EXAMPLE 35

SYNTHESIS OF 4-DIETHYLCARBAMOYLMETHOXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




10 General procedure G for the synthesis of 4-Alkoxy Derivatives.

To a 0.2 M solution of the phenol prepared as in Example 13 (1 eq) in DME or DMF was added alkyl halide and cesium carbonate (Cs₂CO₃). The solution was stirred at ambient temperature until complete, normally 1-12 h. The reaction was diluted with ethyl acetate, 1% HCl was added and the layers separated. The organic layer was washed with 5% NaHCO₃, brine, dried (Na₂SO₄) and concentrated *in vacuo*. Chromatography on SiO₂ afforded the aryl ether.

The title compound was prepared as described in general procedure G employing DME, 1.5 eq of N,N-diethyl-2-chloroacetamide and 2.0 eq of cesium carbonate. Chromatography with 3% ethyl acetate/methylene chloride to 10% ethyl acetate/methylene chloride gradient afforded a 26% yield of product. ESI-MS m/z 554 (MNa⁺).

EXAMPLE 36


SYNTHESIS OF 4-(4-NITRO-BENZYLOXY)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-
 TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 The title compound was prepared as described in general procedure F employing 2.3 eq PPh₃, 2.0 eq. nitrobenzyl alcohol and 2.3 eq DEAD. Chromatography with 10% ethyl acetate/hexane afforded the title compound in 30% yield. ESI-MS m/z 576 (MNa⁺).

EXAMPLE 37


10 SYNTHESIS OF 4-(BIPHENYL-4-YLMETHOXY)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-
 TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as described in general procedure F employing 1.7 eq PPh₃, 1.7 eq. biphenylmethanol and 1.7 eq DEAD. Chromatography with 10% ethyl acetate/hexane afforded the title compound in 47% yield. ESI-MS m/z 607 (MNa⁺).

EXAMPLE 38

SYNTHESIS OF 4-(2-NAPHTHALEN-2-YL-ETHOXY)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL)




5

The title compound was prepared as described in general procedure F employing 2.3 eq PPh₃, 2.0 eq. naphthaleneethanol and 2.3 eq DEAD. Chromatography with 5% ethyl acetate/hexane afforded the title compound in 46% yield. ESI-MS m/z 595 (MNa⁺).

EXAMPLE 39


10 SYNTHESIS OF 4-(3-FLUORO-BENZYLOXY)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as described in general procedure F employing 2.3 eq PPh₃, 2.0 eq. 3-fluorobenzyl alcohol and 2.3 eq DEAD.
15 Chromatography with 5% ethyl acetate/hexane afforded the title compound in 50% yield. ESI-MS m/z 549 (MNa⁺).

EXAMPLE 40


SYNTHESIS OF 11-OXO-4-(3-PHENYL-PROPOXY)-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5 The title compound was prepared as described in general procedure F employing 2.3 eq PPh₃, 2.0 eq. 3-phenyl-1-propanol and 2.3 eq DEAD. Chromatography with 5% ethyl acetate/hexane afforded the title compound in 40% yield. ESI-MS m/z 559 (MNa⁺).

EXAMPLE 41


10 SYNTHESIS OF 11-OXO-4-(2-PYRIDIN-2-YL-ETHOXY)-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



15 The title compound was prepared as described in general procedure F employing 2.3 eq PPh₃, 2.0 eq. 2-(2-hydroxyethyl) pyridine and 2.3 eq DEAD. Chromatography with 20% ethyl acetate/dichloromethane afforded the title compound in 36% yield. ESI-MS m/z 546 (MNa⁺).

EXAMPLE 42


SYNTHESIS OF 4-(2-METHOXY-ETHOXY)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 The title compound was prepared as described in general procedure F employing 2.3 eq PPh₃, 2.0 eq. 2-methoxyethanol and 2.3 eq DEAD. Chromatography with 18% ethyl acetate/hexane, then 15% acetone/hexane afforded the title compound in 50% yield. ESI-MS m/z 499 (MNa⁺), 515 (MK⁺).

EXAMPLE 43


- 10 SYNTHESIS OF 4-CYCLOPENTYLOXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 15 The title compound was prepared as described in general procedure F employing 2.3 eq PPh₃, 2.0 eq. cyclopentanol and 2.3 eq DEAD. Chromatography with 10% ethyl acetate/hexane afforded the title compound in 36% yield. ESI-MS m/z 509 (MNa⁺).

EXAMPLE 44


SYNTHESIS OF 4-(3-CYANO-PROPOXY)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-
 TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 The title compound was prepared as described in general procedure G employing DMF, 1.5 eq 4-bromobutyronitrile and 2.0 eq Cs₂CO₃. Chromatography with 20% ethyl acetate/hexane afforded the title compound in 82% yield. ESI-MS m/z 508 (MNa⁺).

EXAMPLE 45


SYNTHESIS OF 4-(5-METHYL-ISOXAZOL-3-YLMETHOXY)-11-OXO-
 10 TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER
 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- The title compound was prepared as described in general procedure F employing 2.3 eq PPh₃, 2.0 eq. 5-methylisoxazole-3-methanol and 2.3 eq DEAD.
 15 Chromatography with 20% ethyl acetate/dichloromethane afforded the title compound in 36% yield. ESI-MS m/z 536 (MNa⁺).

EXAMPLE 46


SYNTHESIS OF 4-ETHOXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as described in general procedure G employing DMF, 1.5 eq iodoethane and 2.0 eq Cs₂CO₃. Chromatography with 10% ethyl acetate/hexane afforded the title compound in 40% yield. ESI-MS m/z 469 (MNa⁺).

EXAMPLE 47


SYNTHESIS OF 4-METHOXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as described in general procedure G employing DMF, 3.0 eq iodomethane and 2.0 eq Cs₂CO₃. Chromatography with 15% ethyl acetate/hexane afforded the title compound in 54% yield. ESI-MS m/z 455 (MNa⁺).

EXAMPLE 48


SYNTHESIS OF 4-ALLYLOXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5 The title compound was prepared as described in general procedure G employing DMF, 3.0 eq 1,3-diiodopropane and 1.0 eq Cs₂CO₃. Chromatography with 10% ethyl acetate/hexane afforded the title compound in 8% yield. ESI-MS m/z 481 (MNa⁺).

EXAMPLE 49


10 SYNTHESIS OF 11-OXO-4-(PYRIDIN-3-YLMETHOXY)-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



15 The title compound was prepared as described in general procedure G employing DMF, 2.0 eq 3-picolylchloride hydrochloride and 4.0 eq Cs₂CO₃. Chromatography with 40% ethyl acetate/hexane afforded the title compound in 75% yield. ESI-MS m/z 532 (MNa⁺).

EXAMPLE 50


SYNTHESIS OF 11-OXO-4-(PYRIDIN-2-YLMETHOXY)-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 The title compound was prepared as described in general procedure G employing DMF, 2.0 eq 2-picolylchloride.HCl and 4.0 eq Cs₂CO₃. Chromatography with 15% ethyl acetate/hexane afforded the title compound in 72% yield. ESI-MS m/z 532 (MNa⁺).

EXAMPLE 51

- 10 SYNTHESIS OF 4-TERT-BUTOXYCARBONYLMETHOXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}] DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




The title compound was prepared as described in general procedure G employing DMF, 1.1 eq ofttert-butylbromo acetate, and 1.5 eq of Cs₂CO₃.

Chromatography with 20% ethyl acetate/hexane afforded the title compound in 89% yield. ESI-MS m/z 555 (MNa^+).

EXAMPLE 52


5 SYNTHESIS OF 4-(DIMETHOXY-PHOSPHORYLOXY)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as described in general procedure G employing DMF, 1.5 eq dimethylchlorophosphate and 2.0 eq Cs_2CO_3 . Chromatography with 40% ethyl acetate/hexane afforded the title compound in 19% yield. ESI-MS m/z 549 (MNa^+).

EXAMPLE 53

SYNTHESIS OF 11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



A. 11-Oxo-4-trifluoromethanesulfonyloxy-tricyclo[6.2.2.0^{2,7}]dodeca-2(7),3,5-triene-9,10-dicarboxylic acid 10-propyl ester 9-(2-trimethylsilyl-ethyl) ester.


To a 0.3 M solution of the propyl ester of Example 13 in dichloromethane was added DIEA (2 eq) and N-phenyltrifluoromethanesulfonimide (1.1 eq). The reaction was stirred for 18 h at rt, then diluted with dichloromethane, quenched with aqueous ammonium chloride. The phases were partitioned, and the organic layer was separated and washed with 5% aqueous NaHCO₃. The solution was dried (Na₂SO₄), concentrated to dryness and chromatographed with 20% to 30% ethyl acetate/hexane.

B. 11-Oxo-tricyclo[6.2.2.0^{2,7}]dodeca-2(7),3,5-triene-9,10-dicarboxylic acid 10-propyl ester 9-(2-trimethylsilyl-ethyl) ester

A 0.155 M solution of the purified triflate in DMF was treated with Pd(OAc)₂ (0.048 eq), 1,1'-Bis(diphenylphosphino)ferrocene (0.054 eq), triethylamine (7.7 eq) and formic acid (8.5 eq). The reaction was heated to 90°C for 2 days, then diluted with dichloromethane and quenched with aqueous ammonium chloride. The phases were partitioned, and the organic layer was separated, washed with brine, then dried (Na₂SO₄). Chromatography with 30% ethyl acetate/hexane afforded the title compound in 59% yield. ¹H NMR (CDCl₃) 7.27 (dd, 2H), 7.23 – 7.15 (m, 2H), 4.26 (ddd, 2H), 4.04 (d, 1H), 3.96 (ddd, 2H), 3.80 (dd, 1H), 3.69 (dd, 1H), 3.23 (ddd, 1H), 2.43 (dd, 1H), 2.13 (ddd, 1H), 1.57 (dd, 2H), 1.61 – 1.54 (m, 4H), 1.03 (ddd, 2H), 0.88 (t, 3H), 0.04 (s, 9H).

EXAMPLE 54

SYNTHESIS OF 4-HYDROXY-11-(METHYL-HYDRAZONO)-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5

General Procedure H for the Synthesis of 11-Imino Derivatives


To a solution of the product of Example 13 (1 eq, 0.12 – 0.3 M) in methanol is added a selected amino derivative. Sodium acetate may be used as acid scavenger in the case where the nucleophile is added as an acid salt. The reaction is stirred until complete, normally 1-18 h. The crude material is isolated either by concentration *in vacuo* or by extraction using ethyl acetate or diethyl ether.

The title compound was prepared as described in general procedure H using 4 eq. of methylhydrazine. The crude product was isolated by adding diethyl ether and concentrating *in vacuo*. Trituration with diethyl ether afforded a 69% yield of hydrazone as predominately the E isomer. ESI-MS *m/z* 447 (MH⁺), 469 (MNa⁺).

15

EXAMPLE 55

SYNTHESIS OF 4-HYDROXY-11-(PHENYL-HYDRAZONO)-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




20

The title compound was prepared as described in general procedure H using 3 eq. of phenylhydrazine. The crude product was isolated by extraction with ethyl acetate. Trituration with CHCl_3 /hexane afforded a 62% yield of the hydrazone. ESI-MS m/z 509 (MH^+).

5

EXAMPLE 56

SYNTHESIS OF 11-[(2-BROMO-PHENYL)-HYDRAZONO]-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




10

The title compound was prepared as described in general procedure H using 2 eq of 2-bromophenylhydrazine hydrochloride and 2 eq. of sodium acetate. The crude product was isolated by extraction with ethyl acetate. Chromatography on SiO_2 using 20% ethyl acetate/hexane afforded a 61% yield of the hydrazone. ^1H NMR (CDCl_3): 7.48 (dd 1H), 7.38 (dd, 1H), 7.2 (m, 2H), 7.10 (d, 1H), 6.65 (m, 3H), 5.7 (br. S, 1H), 4.25 (m, 3H), 3.97 (m, 2H), 3.80 (m, 1H), 3.63 (m, 1H), 3.20 (m, 1H), 2.50 (dd, 1H), 2.20 (m, 1H), 1.59 (hex, 2H), 1.07 (m, 2H), 0.89 (t, 3H), 0.02 (s, 9H).

15

EXAMPLE 57

SYNTHESIS OF 11-(DIMETHYL-HYDRAZONO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5

The title compound was prepared as described in general procedure H employing 1,1-dimethylhydrazine (1.4 eq). Chromatography using 40% - 50% ethyl acetate/hexane afforded the title compound in 38% yield. ESI-MS m/z 461 (MH^+).

EXAMPLE 58

10 SYNTHESIS OF 4-HYDROXY-11-[(2-HYDROXY-ETHYL)-HYDRAZONO]-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




15 The title compound was prepared as described in general procedure H employing 2-hydroxyethyl hydrazine (1.4 eq) as the reagent. Chromatography using 50% - 80% ethyl acetate/hexane afforded the title compound in 46% yield. ¹H NMR ($CDCl_3$) 7.25 (d, 1H), 6.70 (dd, 1H), 6.65 (d, 1H), 5.3 - 5.1 (br s, 1H), 4.25 (dd, 2H), 4.25 - 4.15

(m, 2H), 3.98 – 3.91 (m, 4H), 3.74 (d, 1H), 3.68 (dd, 1H), 3.21 – 3.19 (m, 1H), 2.41 (dd, 1H), 2.12 (ddd, 1H), 1.58 (dd, 2H), 1.24 (s, 1H), 1.04 (dd, 2H), 0.91 (t, 3H), 0.05 (s, 9H).

EXAMPLE 59


SYNTHESIS OF 11-(THIOSEMICARBAZONO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-
5 2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-
ETHYL) ESTER



The title compound was prepared as described in general procedure H
employing thiosemicarbazide as the reagent. Chromatography using 50% ethyl acetate/hexane
10 afforded the title compound in 42% yield. ESI-MS m/z 492 (MH^+), 514 (MNa^+).

EXAMPLE 60

SYNTHESIS OF 11-(4-METHYL-3-THIOSEMICARBAZONO)-4-HYDROXY-
TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER
9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




15

The title compound was prepared as described in general procedure H employing 4-methyl-3-thiosemicarbazide (1.4 eq) as the reagent. Chromatography using 35% ethyl acetate/hexane afforded the title compound in 35% yield. APCI-MS m/z 506 (MH^+), 505 (M).

5

EXAMPLE 61

SYNTHESIS OF 4-HYDROXY-11-(METHYL-PHENYL-HYDRAZONO)-
TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER
9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




10

The title compound was prepared as in general procedure H using 1.2 eq. of N-methyl-N-phenylhydrazine. The crude product was isolated by concentration *in vacuo*. Chromatography on SiO₂ using 5% ethyl acetate/methylene chloride afforded a 57% yield of the hydrazone. ESI-MS m/z 523 (MH^+), 521 ($M-H$).

EXAMPLE 62


SYNTHESIS OF 11-(METHANESULFONYL-HYDRAZONO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as in general procedure H using 2.4 eq. of methanesulfonyl hydrazine. The crude product was isolated by extraction with ethyl acetate. Chromatography on SiO₂ using 35% ethyl acetate/hexane afforded a 10% yield of the Z-isomer and 50% of the E-isomer. ESI-MS 533.1 (MNa⁺).

EXAMPLE 63


SYNTHESIS OF 11-(BENZENESULFONYL-HYDRAZONO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as in general procedure H using 2.4 eq. of benzenesulfonyl hydrazine. The crude product was isolated by extraction with ethyl acetate. Chromatography on SiO₂ using 30% ethyl acetate/hexane afforded a 62% yield of product. ESI-MS m/z 595 (MNa⁺).

EXAMPLE 64

SYNTHESIS OF 11-(4-METHOXYBENZENESULFONYL-HYDRAZONO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5

The title compound was prepared as described in general procedure H using 2 eq. of 4-methoxybenzenesulfonyl hydrazine. The crude product was isolated by extraction with ethyl acetate. Chromatography on SiO₂ using 40% ethyl acetate/hexane afforded an 82% yield of product. ESI-MS m/z 603 (MH⁺).

10

EXAMPLE 65

SYNTHESIS OF 11-(ACETYL-HYDRAZONO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




15

The title compound was prepared as described in general procedure H using 2.6 eq. of acetyl hydrazine. The crude product was isolated by extraction. Chromatography on SiO₂ using 50% ethyl acetate/hexane afforded a 34% yield of a 4/1 mix of E/Z isomers. ESI-MS m/z 475 (MH⁺).

EXAMPLE 66

SYNTHESIS OF 4-HYDROXY-11-HYDROXYIMINO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5 The title compound was prepared as described in general procedure H employing hydroxylamine (1.1 eq) and sodium acetate (2.3 eq). The crude product was isolated by extraction. Chromatography with 5% Acetone/1% acetic acid/94% dichloromethane followed by recrystallization in dichloromethane/hexane afforded the title compound in 58% yield. ESI-MS m/z 456 (MNa^+).

10

EXAMPLE 67


SYNTHESIS OF 4-HYDROXY-11-METHOXYIMINO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



15 The title compound was prepared as described in general procedure H employing methoxylamine (1.1 eq) and sodium acetate (2.3 eq). The crude product was isolated by extraction. Chromatography with 10% ethyl acetate/dichloromethane resulted in a 34% yield of the less polar isomer: ESI-MS m/z 448 (MNa^+), 470 (MNa^+) and a 34% yield of the more polar isomer: ESI-MS m/z 448 (MH^+), 470 (MNa^+).

EXAMPLE 68

SYNTHESIS OF 4-HYDROXY-11-PHENOXYIMINO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5 The title compound was prepared as described in the general procedure H employing 1.1 eq of O-phenylhydroxylamine and 2.3 eq of sodium acetate. The crude product was isolated by extraction. Chromatography with 30% ethyl acetate/dichloromethane afforded the title compounds as a 1:1 mixture in 63% combined yield. ESI-MS m/z 510 (MH^+), 532 (MNa^+).

10

EXAMPLE 69


SYNTHESIS OF 11-BENZYLOXYIMINO-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



15 The title compound was prepared as described in general procedure H employing 1.1 eq of benzylhydroxylamine and 2.3 eq of sodium acetate. The crude product was isolated by extraction. Chromatography with 30% ethyl acetate/dichloromethane afforded the title compounds as a 2:1 mixture of Z and E isomers in 47% combined yield. ESI-MS m/z 524 (MH^+).

EXAMPLE 70

SYNTHESIS OF 4-HYDROXY-11-(4-NITRO-BENZYLOXYIMINO)-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5

The title compound was prepared as described in general procedure H employing 1.1 eq of (4-nitrobenzyl)hydroxylamine and 2.3 eq of sodium acetate. The crude product was isolated by extraction. Chromatography with 20–30% ethyl acetate/dichloromethane afforded the title compound in 28% yield. ESI-MS m/z 569 (MH^+).

10

EXAMPLE 71

SYNTHESIS OF 11-(5-CHLORO-[1,2,3]THIADIAZOL-4-YLMETHOXYIMINO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




15

The title compound was prepared as described in general procedure H employing 1.1 eq of (4-chloro)thiadiazolyl-5-methoxyhydroxylamine and 2.3 eq of sodium acetate. The crude product was isolated by extraction. Chromatography with 30% - 40%

ethyl acetate/dichloromethane afforded the title compounds as a 1:1 mixture of E and Z isomers in 92% yield. ESI-MS m/z 566 (MNa^+).

EXAMPLE 72


SYNTHESIS OF 11-(3-FLUORO-BENZYLOXYIMINO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-
5 2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as described in general procedure H employing 1.1 eq of (3-fluoro)benzylhydroxylamine and 2.3 eq of sodium acetate. The
10 crude product was isolated by extraction. Chromatography with 30% ethyl acetate/dichloromethane afforded the title compounds as a 1:1.3 mixture of E and Z isomers in 89% yield. ESI-MS m/z 542 (MNa^+).

EXAMPLE 73


SYNTHESIS OF 4-HYDROXY-11-[2-OXO-2-(4-PHENYL-PIPERAZIN-1-YL)-ETHOXYIMINO]-
15 TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was prepared as described in general procedure H employing 2-aminooxy-1-(4-phenyl-piperazin-1-yl)-ethanone (1.1 eq) and sodium acetate (2.3 eq). The crude product was isolated by extraction. Chromatography with 40% - 60% ethyl acetate/dichloromethane afforded the less polar E isomer in 8% yield and the more polar Z isomer in 18% yield. ESI-MS m/z 636 (MH^+), 658 (MNa^+).

EXAMPLE 74

SYNTHESIS OF 11-(4-FLUORO-BENZYLOXYIMINO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




10

The title compound was prepared as described in general procedure H employing (4-fluoro)benzylhydroxylamine (1.1 eq) and sodium acetate (2.3 eq). The crude product was isolated by extraction. Chromatography with 0% - 10% ethyl acetate/dichloromethane afforded the title compound in 25% yield. ¹H NMR (CDCl₃) 7.32 (dd, 2H), 7.09 – 6.99 (m, 3H), 6.69 – 6.64 (m, 2H), 5.05 – 5.03 (m, 3H), 4.95 (s, 1H), 4.24 (dd, 2H), 3.95 (ddd, 2H), 3.64 (d, 1H), 3.50 (dd, 1H), 3.08 – 3.06 (m, 1H), 2.50 (dd, 1H), 2.12 (ddd, 1H), 1.58 (dd, 2H), 1.02 (dd, 2H), 0.87 (t, 3H), 0.04 (s, 9H).

EXAMPLE 75

SYNTHESIS OF 4-HYDROXY-11-(2-PHENOXY-ETHOXYIMINO)-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5

The title compound was prepared as described in general procedure H employing (3-phenoxy)ethylhydroxylamine (1.1 eq) and sodium acetate (2.3 eq). The crude product was isolated by extraction. Chromatography with 0% - 10% ethyl acetate/dichloromethane afforded the title compound in 22% yield. ¹H NMR (CDCl₃) 7.31 – 7.30 (m, 2H), 7.06 (d, 1H), 6.97 – 6.92 (m, 3H), 6.64 (dd, 1H), 6.57 (d, 1H), 5.03 (d, 1H), 4.87 (s, 1H), 4.41 – 4.37 (m, 2H), 4.25 (ddd, 2H), 4.16 (dd, 2H), 3.94 (ddd, 2H), 3.63 (dd, 1H), 3.51 (dd, 1H), 3.09 – 3.06 (m, 1H), 2.52 (dd, 1H), 2.13 (ddd, 1H), 1.57 (ddd, 2H), 1.03 (dd, 2H), 0.86 (t, 3H), 0.05 (s, 9H).

EXAMPLE 76

15 SYNTHESIS OF 11-ALLYLOXYIMINO-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




The title compound was prepared as described in general procedure H employing O-allylhydroxylamine (1.1 eq) and sodium acetate (2.3 eq). The crude product was isolated by extraction. Chromatography with 25% - 40% ethyl acetate/hexane afforded the title compound in 45% yield. ESI-MS m/z 496 (MNa^+).

5

EXAMPLE 77

SYNTHESIS OF 11-(2,4-DICHLOROBENZYL-OXIMO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




10

The title compound was prepared as described in general procedure H using 1.1 eq of O-2,4-dichlorobenzyl hydroxyl amine hydrochloride and 1.1 eq of sodium acetate. The crude product was isolated by concentrating *in vacuo*. Chromatography on SiO_2 using 25% ethyl acetate/hexane followed by 3% ethyl acetate/methylene chloride afforded a 28% yield of Z-isomer, the less polar compound, and 33% yield of the E-isomer, the more polar compound. ESI-MS m/z , Z-isomer 592 (MH^+), 594 ($(M + 2)H^+$); E-isomer 592 (MH^+), 594 ($(M + 2)H^+$).

15

EXAMPLE 78


SYNTHESIS OF 11-(SEMICARBAZONO)-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 The title compound was prepared as described in general procedure H using 1.3 eq of semicarbazide hydrochloride and 1.3 eq of sodium acetate. The crude product was isolated by extraction with ethyl acetate. Chromatography on SiO₂ using 70% ethyl acetate/hexane afforded a 48% yield of product. ESI-MS *m/z* 498 (MNa⁺).

EXAMPLE 79


- 10 SYNTHESIS OF (9,10 TRANS)-10-ALLYLOXYCARBONYLAMINO-4,11-DIHYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- To a solution of the ketone prepared in Example 25 (12.4 mg, 0.029 mmol) in 0.5 mL methanol was added sodium borohydride (21.1 mg, 0.56 mmol). After 20 min
15 H₂O was added, the solution acidified to pH 1 with 1% HCl, and the product extracted with ethyl acetate. The organic layer was washed with H₂O, brine, dried (Na₂SO₄) and concentrated. Flash chromatography (SiO₂, 45% ethyl acetate/dichloromethane) afforded 3.7 mg (30%) of the less polar alcohol and 4.1 mg (33%) of the more polar alcohol. ESI-MS *m/z*: less polar product 456 (MNa⁺), more polar product 456 (MNa⁺).

EXAMPLE 80


SYNTHESIS OF (9,10 CIS)-10-ALLYLOXYCARBONYLAMINO-4,11-DIHYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 To a solution of the ketone from Example 3 (25.9 mg, 0.06 mmol) in methanol (1.1 mL) held at 20°C with a water bath was added sodium borohydride (44.7 mg, 1.2 mmol). The reaction was stirred for 20 min, diluted with ethyl acetate and quenched with water followed by 1% HCl. The layers were separated, and the organic layer washed with 5% NaHCO₃, water, brine, dried (Na₂SO₄), and concentrated *in vacuo*.
- 10 Chromatography on SiO₂ using 40% ethyl acetate/hexane afforded 13.2 mg (50%) of product as the less polar diastereomer. ESI-MS *m/z* 456 (MNa⁺).

EXAMPLE 81

SYNTHESIS OF 4,11-DIHYDROXY-11-PHENYL-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



15


To a solution of the compound of Example 13 (61 mg, 0.146 mmol) in tetrahydrofuran (2.9 mL) at -10° was added phenylmagnesium bromide (360 µL, 1 M).

Additional aliquots (720 μ L) of phenylmagnesium bromide were added at 30 min intervals until the reaction was complete by TLC. The cooling bath was removed, the reaction diluted with ethyl acetate, then quenched with 3% HCl. The layers were separated and the organic layer washed with 1% HCl, 5% NaHCO₃, H₂O, brine, and dried (Na₂SO₄).

- 5 Concentration *in vacuo* followed by chromatography afforded 26 mg (36%) of the tertiary alcohol. ESI-MS m/z 519 (MNa⁺).

EXAMPLE 82

SYNTHESIS OF 4-HYDROXY-11-PROPYLAMINO-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




10

- To a solution of the compound of Example 13 (75 mg, 0.18 mmol) in methanol (1 mL) was added n-propylamine (75 μ L, 1.8 mmol) and acetic acid (52 μ L, 1.8 mmol). After 10 min sodium triacetoxy borohydride (380 mg, 1.8 mmol) was added and the solution stirred overnight. Additional aliquots of n-propylamine (600 μ L), acetic acid (500 μ L) and sodium triacetoxy borohydride (400 mg) were added and the reaction allowed to proceed for 1.5 h. The reaction was diluted with ethyl acetate and 5% NaHCO₃. The layers were separated, and the organic layer washed with 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*. Chromatography on SiO₂ using a gradient of 45% ethyl acetate/methylene chloride - 2% methanol/methylene chloride - 8% methanol/methylene chloride afforded a 34% yield of the less polar amine and 37% yield of the more polar amine.
- 15
- 20 ESI-MS m/z 462 (MH⁺).

EXAMPLE 83

SYNTHESIS OF 4-HYDROXY-11-(4-METHYL-BENZYLAMINO)-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




5

To a solution of the compound of Example 13 (50 mg, 0.12 mmol) in methanol (0.5 mL) was added acetic acid (14 μ L, 0.24 mmol), 4-methylbenzyl amine (31 μ L, 2.1 eq.) and sodium cyanoborohydride (38 mg, 0.60 mmol). After 2 h an additional aliquot of sodium cyanoborohydride (10 mg) was added and the reaction allowed to stir for 30 more min. The reaction was then quenched with 3% HCl and diluted with ethyl acetate. The biphasic mixture was then basified to pH 8 with 5% NaHCO₃. The layers were separated, and the organic layer washed with H₂O, brine, dried (Na₂SO₄) and concentrated *in vacuo*. Chromatography on SiO₂ using first a gradient of 15% ethyl acetate/hexane - 25% ethyl acetate/hexane followed by a second chromatography using a gradient 10% ethyl acetate/methylene chloride - 17% ethyl acetate/methylene chloride afforded a 19% yield of the less polar amine and 16% of the more polar amine. ESI-MS *m/z* 524 (MH⁺).

15

EXAMPLE 84


SYNTHESIS OF 4-HYDROXY-11-METHYLAMINO-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



20

EXAMPLE 86


SYNTHESIS OF 11-DIMETHYLAMINO-4-HYDROXY-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 To a solution of the compound of Example 13 (79.5 mg, 0.19 mmol) in methanol (0.5 mL) was added dimethylamine (40 wt% in H₂O, 600 μ L, 4.8 mmol) and acetic acid (330 μ L, 5.7 mmol). After 10 min sodium triacetoxy borohydride (1.0g, 4.8 mmol) was added. Additional aliquots of reagents were added at 3 h and 4.5 h, and the reaction allowed to proceed for 2 h after the final addition of reagents. The reaction was
- 10 quenched with H₂O and diluted with ethyl acetate. The layers were separated and the organic layer washed with 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄) and concentrated. *in vacuo*. Flash chromatography on SiO₂ with 5% methanol/methylene chloride afforded 9.4 mg of a single diastereomer.

EXAMPLE 87


- 15 SYNTHESIS OF 11-[ACETYL-(4-METHYL-BENZYL)-AMINO]-4-ACETOXY-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



To a solution of the more polar amine from Example 83 (16.5 mg, 0.03 mmol) in methylene chloride (0.4 mL) was added N-methylmorpholine (12 μ L, 0.10 mmol) followed by acetic anhydride (10 μ L, 0.10 mmol). The reaction was stirred for 16 h at room temp, diluted with ethyl acetate, and quenched with 5% NaHCO₃. The layers were separated, and the organic layer was washed with H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*. Flash chromatography on SiO₂ with 40% ethyl acetate/hexanes afforded the amide. ESI-MS m/z 608 (MH⁺), 630 (MNa⁺).

EXAMPLE 88

SYNTHESIS OF 11-[ACETYL-METHYLAMINO]-4-ACETOXY-TRICYCLO[[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



A. Diastereomer 1.


To a solution of the less polar amine from Example 84 (35 mg, 0.08 mmol) in dichloromethane (1.1 mL) was added NMM (27 μ L, 0.24 mmol) and acetic anhydride (15.3 μ L, 0.16 mmol). The reaction was stirred for 18 h at ambient temperature, diluted with ethyl acetate and quenched with 1% HCl. The layers were separated, and the organic layer washed with 1% HCl, 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄) and concentrated *in vacuo*. Chromatography on SiO₂ using a gradient of 45% ethyl acetate/hexane – 55% ethyl acetate/hexane afforded 9.3 mg (22%) of the bis-acylated product, which by TLC is slightly less polar than the free phenol product. ¹H NMR (CDCl₃, 53°C): 7.19 (d, 1H), 6.95 (m, 2H), 4.30 (m, 2H), 3.89 (t, 2H), 3.54 (m, 3H), 3.20 (s, 3H), 2.96 (d, 1H), 2.26 (s, 3H), 2.13 (s, 3H), 1.80 (m, 2H), 1.59 (m, 2H), 1.06 (sextet, 2H), 0.91 (t, 2H), 0.07 (s, 9H).

B. Diastereomer 2.

To a solution of the more polar amine from Example 84 (58 mg, 0.13 mmol) in methylene chloride (2 mL) was added N-methylmorpholine (43 μ L, 0.39 mmol) and acetic anhydride (15 μ L, 0.16 mmol). The reaction was stirred at room temp for 21 h, diluted with ethyl acetate, and quenched with 1% HCl. The layers were separated, and the organic layer washed with 1% HCl, 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*. Multiple flash chromatographies on SiO₂ using 2% methanol/methylene chloride afforded 34 mg (53%) of the phenol as the more polar product and 14.7 mg (21%) of the phenol acetate as the less polar product. ¹H NMR (CDCl₃, 21°C, ca. 3:1 mix of rotamers): 7.24 (d, 1H), 6.90 (m, 2H), 5.21 (m, 0.75H), 4.23 (m, 2.25H), 3.90 (t, 2H), 3.52 (m, 3H), 2.99 (m, 1H), 2.30-1.96 (m, 10H), 1.56 (sextet, 2H), 1.30 (m, 1H), 1.03 (m, 2H), 0.93 (t, 2H), 0.05 (s, 9H).

EXAMPLE 89


SYNTHESIS OF 11-[ACETYL-METHYLAMINO]-4-HYDROXY-TRICYCLO[[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



The title compound was isolated from the preparation of Diastereomer 2 of Example 88. By TLC it is more polar than the corresponding phenol acetate. ESI-MS *m/z* 476 (MH⁺), 498 (MNa⁺), 474 (M-H)⁻.

EXAMPLE 90

SYNTHESIS OF 4,11-DIHYDROXY-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-(2,4-DIMETHOXY-BENZYL) ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



5


To a solution of the ketone of Example 9 (31.1 mg, 0.06 mmol) in methanol (1 mL) at 15°C was added NaBH₄ (44.0 mg, 1.16 mmol). The reaction was stirred for 45 min, diluted with ethyl acetate, and quenched with 1% HCl. The layers were separated and the organic layer washed with 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*. Flash chromatography on SiO₂ using 30% ethyl acetate/methylene chloride afforded 6.5 mg (21%) of the less polar alcohol and 13.8 mg (44%) of the more polar alcohol. ESI-MS *m/z*, less polar product 551 (MNa⁺), more polar product 551 (MNa⁺).

10

EXAMPLE 91

SYNTHESIS OF 11-SPIRO-(1,4-DIOXACYCLOPENTYL)-4-HYDROXY-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER

15




To a solution of the ketone of Example 13 (56.0 mg, 0.133 mmol) in benzene (1.3 mL) was added ethylene glycol (45 µL) and *p*-tuenesulfonic acid monohydrate (4.5

mg). The solution was refluxed for 1 h, using a Dean-Stark trap to collect the water. Upon cooling the reaction was diluted with ethyl acetate and 5% NaHCO₃ was added. The layers were separated, and the organic layer washed with 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*. Flash chromatography on SiO₂ using 25% ethyl acetate/hexane
5 afforded 14.1 mg (23%) of the ketal. ESI-MS m/z 485 (MNa⁺).

EXAMPLE 92

SYNTHESIS OF 11-ETHOXYCARBONYLMETHYLENE-4-HYDROXY- TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




10

To a solution of triethylphosphonoacetate (59 μ L, 0.30 mmol) in tetrahydrofuran (0.5 mL) at 0°C was added a solution of potassium hexamethyldisilazide (0.45 M in toluene, 0.685 mL, 0.31 mmol). After stirring for 15 min at 0°C a solution of the ketone from Example 13 (59.6 mg, 0.14 mmol) in tetrahydrofuran (0.9 mL) was added. The
15 reaction was stirred for 30 min at 0°C, 3 h at ambient, then placed in a refrigerator without stirring for 66 h. Upon removal, the reaction was stirred at ambient temperature for 6 h, diluted with ethyl acetate, and quenched with 1% HCl. The layers were separated, and the organic layer washed with 1% HCl, 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*. Flash chromatography on SiO₂ using 17% ethyl acetate/hexane
20 afforded 32.2 mg (46%) of product as a 1/1 mixture of E/Z isomers. ESI-MS m/z 511 (MNa⁺).

EXAMPLE 93


SYNTHESIS OF 4-HYDROXY-11-METHYLENE-TRICYCLO[[6.2.2.0^{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- 5 To a solution of methyl triphenylphosphonium bromide (236 mg, 0.66 mmol) in tetrahydrofuran (0.5 mL) was added potassium hexamethyldisilazide (0.5 M in toluene, 1.3 mL, 0.65 mmol). The yellow-orange solution was stirred for 15 min at ambient temp, then a solution of the ketone from example 13 (63.5 mg, 0.15 mmol) in tetrahydrofuran (0.3 mL) was added. After 30 min at room temp the reaction was diluted
- 10 with ethyl acetate and quenched with H₂O. The layers were separated, and the organic layer washed with 1% HCl, 5% NaHCO₃, H₂O, brine, dried (Na₂SO₄), and concentrated *in vacuo*. Flash chromatography on SiO₂ using 15% ethyl acetate/hexane afforded 34.2 mg (54%) of the olefin. ¹H NMR (CDCl₃), 7.05 (2H), 6.63 (2H), 5.10 (1H), 4.96 (1H), 4.73 (1H), 4.25 (2H), 3.95 (3H), 3.55 (2H), 3.07 (1H), 2.49 (1H), 2.08 (1H), 1.60 (2H), 1.06
- 15 (2H), 0.88 (3H), 0.09 (9H).

EXAMPLE 94


SYNTHESIS OF 4,11-DIHYDROXY-TRICYCLO[6.2.2.0_{2,7}]]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



To a 0.1 M solution of ketone from Example 13 in methanol was added NaBH₄ (10 eq). After stirring at rt for 20 min, the reaction was quenched with saturated aqueous ammonium chloride and diluted with dichloromethane. The aqueous phase was acidified with 1 M HCl, the phases were partitioned, and the aqueous phase extracted 3 x
5 with dichloromethane. Organic extracts were combined, dried (Na₂SO₄) and concentrated. Chromatography with 40% ethyl acetate/hexane afforded the less polar isomer in 40% yield ESI-MS m/z 419 (M-H)⁻, and the more polar isomer in 20% yield ESI-MS m/z 419 (M-H)⁻.

EXAMPLE 95

10 SYNTHESIS OF 11-AMINO-4-HYDROXY-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




To a 0.1 M solution of ketone from Example 13 in methanol was added 4-methylbenzylamine (3 eq), Na(OAc)₃BH (3 eq), acetic acid (3 eq) and 3A mol. sieves (300 mg/mL methanol). The reaction was stirred overnight, then quenched with aqueous
15 NaHCO₃, and extracted with ethyl acetate. The organic extracts were combined, dried (Na₂SO₄) and concentrated. Chromatography with 5% - 30% ethyl acetate/hexane afforded the less polar isomer in 29% yield.

The product from the preceding reaction was dissolved in enough ethanol to give a 0.07 M solution, which was treated with 20% palladium hydroxide on carbon
20 (135 mg/mmol st. mat) and acetic acid (24 eq.). The reaction was stirred at rt under an atmosphere (balloon) of hydrogen gas. After 1.5 h, the reaction was filtered over Celite with dichloromethane and concentrated. Chromatography using 10% - 20%

methanol/dichloromethane afforded a single isomer in 45% yield. ESI-MS m/z 420 (MH^+), 418 ($M-H$).

EXAMPLE 96


5 SYNTHESIS OF 4,11-DIHYDROXY-11-METHYL-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-
TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



To a 0°C solution of ketone from Example 13 in THF (0.12 M) was added methylmagnesium bromide (1.4 M in THF/toluene, 6.7 eq). After complete reaction and aqueous workup, the crude product was purified by column chromatography using 25% -
10 40% ethyl acetate/hexane. The less polar isomer was isolated in 12% yield. ¹H NMR (CDCl₃) 7.26 (d, 1H), 6.99 (s, 1H), 6.82 (dd, 1H), 4.94 (s, 1H), 4.27 (ddd, 2H), 3.90 (dd, 2H), 3.90 – 3.85 (m, 1H), 3.35 (d, 1H), 3.28 (d, 1H), 3.18 – 3.15 (m, 1H), 1.77 (dd, 1H), 1.57 (dd, 2H), 1.44 – 1.39 (m, 1H), 1.25 (s, 2H), 1.06 (ddd, 2H), 1.01 (s, 3H), 0.89 (dd, 3H), 0.06 (s, 9H).

EXAMPLE 97

15 SYNTHESIS OF 4,11-DIHYDROXY-11-METHYL-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-
TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



Cerium chloride (2.0 eq) was heated under vacuum, cooled and suspended in THF. The 0.3 M solution was cooled to -75°C and treated with methylmagnesium


bromide (1.4 M in THF/toluene, 4 eq) in a dropwise fashion. The slurry was stirred for 1.5 h, at which point the ketone from Example 13 was added as a 0.3 M solution in THF. The reaction was stirred at -75°C for 2 h, then warmed to rt. After 1.5 h, reaction quenched with saturated aqueous ammonium chloride and extracted with ethyl acetate.

- 5 Chromatography using 30% - 40% ethyl acetate/hexane afforded the more polar isomer in 13% yield. ¹H NMR (CDCl₃) 7.26 (d, 1H), 6.70 (s, 1H), 6.68 (dd, 1H), 5.26 (s, 1H), 4.25 (ddd, 2H), 3.93 (ddd, 2H), 3.57 (dd, 1H), 3.42 - 3.40 (s, 1H), 3.31 - 3.29 (s, 1H), 2.92 - 2.91 (s, 1H), 1.72 (dd, 1H), 1.57 (dd, 2H), 1.56 (s, 3H), 1.40 (dm, 1H), 1.26 (s, 1H), 1.04 (dd, 2H), 0.87 (t, 3H), 0.06 (s, 9H).

10

EXAMPLE 98

SYNTHESIS OF 4,11-DIHYDROXY-11-HYDROXYMETHYL-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER




15

The olefin of Example 93 (61077) was dissolved in enough 1:1 tert-butanol:water to give a 0.06 M solution. The solution was treated with OsO₄ (4 wt.% soln. in water, 0.03 eq) and 4-methyl morpholine-N-oxide (3 eq) and heated to 50°C. After stirring overnight, the reaction was quenched with sodium bisulfite. Celite was added, and the solution allowed to stir an additional 3 hours. The solution was then diluted 20 fold with THF and filtered over a short plug of silica. The crude solids were purified by column chromatography (50% - 60% ethyl acetate/hexane) to afford the less polar isomer in 36% yield. APCI-MS m/z 449 (M-H)⁻.

20

EXAMPLE 99

SYNTHESIS OF 10-(BENZYL-METHYL-CARBAMOYL)-5-HYDROXY-12-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-CARBOXYLIC ACID ALLYL ESTER




5 General Procedure I for the Synthesis of 5-Hydroxy-10-amido Derivatives

A solution of the trans allyl TMS-ethyl ester prepared in Example 19 (1 equivalent) in 2 mL of TFA/H₂O (95%) was allowed to stir at rt for 30 min. The volatiles were evaporated, acetonitrile (2 mL) and toluene (5 mL) were added and the resulting solution was concentrated to dryness (2x) to afford the crude carboxylic acid. The white residue was dissolved in DMF (0.6 mL) and a selected amine (1.5 equivalents), HATU (1.2 equivalents) and NMM (2.7 equivalents) were added and the reaction mixture was allowed to stir at rt under nitrogen overnight. The solution was concentrated to dryness and dichloromethane or ethyl acetate was added (10 mL) and the organic layer was washed with HCl (1 N, 3x10 mL), NaHCO₃ solution (5%, 2x10 mL) and brine (1x10 mL). Upon drying (MgSO₄) the organic layer, the filtered solution was concentrated to dryness and column chromatography provided the desired product.

The title compound was prepared as described in general procedure I using N-methylbenzylamine, resulting in a yield of 19% (10 mg). ESI-MS m/z 442 (MNa⁺), 418 (M-H)⁻.

EXAMPLE 100


SYNTHESIS OF 5-HYDROXY-12-OXO-10-PROPYLCARBAMOYL-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9-CARBOXYLIC ACID ALLYL ESTER



5 The title compound was prepared as described in general procedure I using propylamine, resulting in a yield of 55% (24 mg). ESI 380 (MNa⁺), 356 (M-H).

EXAMPLE 101

SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-[2-(TOLUENE-4-SULFONYL)-ETHYL] ESTER



10


General Procedure J for the Synthesis of 9-Esters

15 To a 0.5 M solution of the corresponding 9-carboxylic acid prepared as described in General Procedure I in 5% DMF/THF was added TSTU (2 eq), 4-methylmorpholine (4 eq), DMAP (2 eq) and a selected alcohol (2 eq). The reaction was allowed to proceed overnight at ambient temperature, after which the reaction was quenched with saturated aqueous ammonium chloride and diluted with 1:1 ethyl acetate:hexane. Aqueous further acidified (pH ~ 2) with 1 M HCl (aq). The phases were partitioned, and the organic layer separated and washed with brine. Solution dried (Na₂SO₄) and concentrated. The products were purified by column chromatography.

Reaction run as in general procedure J using 2-(p-Tosyl)ethanol. Chromatography (ethyl acetate/hexane) affords the title compound in 61% yield. ESI-MS m/z 523 (MNa^+), 499 ($M-H$)⁻.

EXAMPLE 102


- 5 SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 9-(3,3-DIMETHYL-BUTYL) ESTER 10-PROPYL ESTER



- Reaction run as in general procedure J using 3,3 dimethylbutanol. Chromatography (30% ethyl acetate/hexane) affords the title compound in 55% yield. ESI-MS m/z 425 (MNa^+).
- 10

EXAMPLE 103


- SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 9-(2-ADAMANTAN-1-YL-ETHYL) ESTER 10-PROPYL ESTER



- Reaction run as in general procedure J using 1-adamantaneethanol. Chromatography (ethyl acetate/hexane) affords the title compound in 87% yield. ESI-MS m/z 479 ($M-H$)⁻.
- 15

EXAMPLE 104


SYNTHESIS OF 4-HYDROXY-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2(7),3,5-TRIENE-9,10-DICARBOXYLIC ACID 10-PROPYL ESTER 9-(3-TRIMETHYLSILANYL-PROPYL) ESTER



- 5 Reaction run as in general procedure J using 1-(trimethylsilyl)-3-propanol. Chromatography (25% - 30% ethyl acetate/hexane) affords the title compound in 38% yield. APCI-MS m/z 433 (MH^+), 431 ($M-H$).

EXAMPLE 105

SYNTHESIS OF 4-(4-CARBOXYMETHOXY-BENZYLOXY)-11-OXO-TRICYCLO[6.2.2.0^{2,7}]DODECA-2,4,6-TRIENE-9,10-DICARBOXYLIC ACID 10-ALLYL ESTER 9-(2-TRIMETHYLSILANYL-ETHYL) ESTER



- A. Butyldimethylsilyl [4-(tert-butyl-dimethyl-silanyloxymethyl)-phenoxy]-acetate:
 To a solution of (4-hydroxymethylphenoxy)acetic acid (10 g, 55 mmol) in
 15 dichloromethane (200 mL) was added *t*-butyldimethylsilyl chloride (18.2 g, 121 mmol), diisopropylethylamine (24 mL, 17.8 g, 138 mmol) and dimethylaminopyridine (2.7 g, 22

mmol). The resulting reaction mixture was allowed to stir at rt for 1.5 h, after which time it was diluted with an additional 300 mL of dichloromethane and washed with 0.1 M citric acid (2x300 mL) and brine (3x300 mL). The resulting yellow organic layer was dried (MgSO₄), filtered and concentrated to dryness to give a white solid, wt. 22.5 g (quantitative). ¹H NMR (CDCl₃) 7.23 (d, *J*=8.8 Hz, 2H), 6.85 (d, *J*=8.8 Hz, 2H), 4.67 (s, 2H), 4.58 (s, 2H), 0.92 (s, 9H), 0.89 (s, 9H), 0.28 (s, 6H), 0.08 (s, 6H).

B. Synthesis of 2-(Toluene-4-sulfonyl)-ethyl (4-hydroxymethyl-phenoxy)-acetate:

a) To a solution of the product from Step A (22.5 g, 55 mmol) in dichloromethane (200 mL) was added 2-toluenesulfonylethanol (26.4 g, 132 mmol), HATU (25 g, 65.8 mmol) and diisopropylethylamine (23 mL, 17.1 g, 132 mmol). The resulting reaction mixture was allowed to stir at rt overnight under nitrogen atmosphere, after which time it was concentrated to dryness, diluted with ethyl acetate (400 mL), washed with 0.1 M citric acid (3x330 mL), 5% NaHCO₃ solution (2x100 mL) and brine (2x100 mL). The resulting organic layer was dried (MgSO₄), filtered and concentrated to dryness to give a brown solid, wt. 49.41 g (185%). Used as is without further purification.

b) The brown residue obtained above was suspended in 80% acetic acid/water solution (500 mL) and allowed to stir at rt for 3 h. The resulting cloudy solution was then concentrated to dryness and used dichloromethane/toluene mixture to get rid of residual acetic acid. Column chromatography (45% acetone/hexane) provided the desired product, which upon trituration with methanol provided the desired product as a white solid, wt. 15.6 g (78% for the three steps). ¹H NMR (CDCl₃) 7.78 (dt, *J*=8.4, 1.9 Hz, 2H), 7.36 (d, *J*=8.0 Hz, 2H), 7.27 (dt, *J*=9.2, 2.5 Hz, 2H), 6.80 (dt, *J*=9.2, 2.5 Hz, 2H), 4.60 (s, 2H), 4.51 (t, *J*=6.1 Hz, 2H), 4.39 (s, 2H), 3.45 (t, *J*=6.0 Hz, 2H), 2.41 (s, 3H). ESI-MS *m/z* 387 (MNa⁺).

C. 4-(4-Carboxymethoxy-benzyloxy)-11-oxo-tricyclo[6.2.2.0^{2,7}]dodeca-2,4,6-triene-9,10-dicarboxylic acid 10-allyl ester 9-(2-trimethylsilanyl-ethyl) ester


a) To a solution of the phenol of Example 19 (17.4 g, 42 mmol), 2-(toluene-4-sulfonyl)ethyl (4-hydroxymethylphenoxy)acetate (18.2 g, 50 mmol) and

triphenylphosphine (13.1g, 50 mmol) in anhydrous tetrahydrofuran (175 mL) was added DEAD (8.0 mL, 8.85g, 51 mmol). The resulting reaction mixture was allowed to stir at rt for 5 h, after which time it was concentrated to dryness. The residue was then dissolved in ethyl acetate (1.0 L) and washed with 0.1 M citric acid (2x100 mL), 5% NaHCO₃ solution (2x100 mL) and brine (2x100 mL). The resulting organic layer was dried (MgSO₄), filtered and concentrated to dryness to give a yellow oil. Column chromatography (4% acetonitrile/dichloromethane) provided the desired tosylethyl ester, wt. 23.9 g (75%).

b) To a solution of the above ester (23.9 g, 31 mmol) in acetonitrile (200 mL) was added piperidine (7.5 mL, 6.5 g, 76 mmol) and DBU (5.6 mL, 5.7 g, 37.4 mmol). The resulting mixture was allowed to stir at rt for 45 min, after which the solution was concentrated to dryness and redissolved in ethyl acetate (1 L). The organic layer was washed with 0.1 N HCl solution (850/150/150 mL) and brine (2x150 mL), dried (MgSO₄), filtered and the solvent was evaporated to give a yellow oil. Column chromatography (2% methanol/dichloromethane, 2 L, followed by 2% methanol/dichloromethane with 2% AcOH, 3 L) provided the desired product as a foamy off-white solid, wt. 15.2 g (63% over two steps). ¹H NMR (CDCl₃) 7.35 (d, *J*=8.5 Hz, 2H), 7.19 (d, *J*=8.2 Hz, 1H), 6.94 (d, *J*=8.8 Hz, 2H), 6.85 (dd, *J*=8.1, 2.6 Hz, 1H), 6.80 (d, *J*=2.5 Hz, 1H), 5.88-5.75 (m, 1H), 5.28-5.19 (m, 2H), 4.94 (s, 2H), 4.69 (s, 2H), 4.49 (dq, *J*=5.6, 1.3 Hz, 2H), 4.30-4.27 (m, 2H), 4.01 (d, *J*=2.5 Hz, 1H), 3.77 (q, *J*=2.7 Hz, 1H), 3.73 (dd, *J*=5.9, 2.3 Hz, 1H), 3.23 (dt, *J*=5.8, 2.3 Hz, 1H), 2.42 (dd, *J*=19.0, 2.2 Hz, 1H), 2.13 (dt, *J*=18.7, 2.6 Hz, 1H), 1.07-1.01 (m, 2H), 0.06 (s, 9H). APCI-MS 603.3 (MNa⁺), 579.2 (M-H).

EXAMPLE 106

PROCEDURE FOR THE SYNTHESIS OF A LIBRARY OF REPRESENTATIVE BENZOBICYCLOOCTANES



5 A. Loading scaffold onto TentaGel Amine Resin (Novabiochem A18764):

1. Place dry resin (5 g, 0.43 mmol/g loading) in Schlenk ware
2. Swell resin with dichloromethane (2x30 mL, 2 min)
3. Add NMP, bubble N₂ through frit and drain (4x30 mL, 5 min)
4. Add the following solutions to swelled resin: NMP solution of Scaffold (0.5 M, 6 mL), NMP solution of DIEA (1.25 M, 7 mL) and NMP solution of HATU (0.5 M, 6.5 mL)
5. Bubble N₂ through frit for 1.5 h; drain
6. Wash resin with NMP (3x30 mL, 5 min)
7. Kaiser test of resin proved negative (Kaiser, E. et al, *Anal. Biochem.*, 1970, 34, 595)
8. Add the following solutions to swelled resin: NMP solution of Scaffold (0.5 M, 4 mL), NMP solution of DIEA (1.25 M, 4 mL) and NMP solution of HATU (0.5 M, 4 mL)
9. Repeat steps 5 - 7
10. Wash resin with 1:1 NMP/dichloromethane (2x30 mL, 3 min)
11. Wash resin with 1:4 NMP/dichloromethane (2x30 mL, 3 min)

12. Wash resin with neat dichloromethane (3x30 mL, 3 min)
 13. Dry under vacuum overnight
 14. Distribute resin into 96-well plate (50 mg/well); store at about -15/-20°C in a ziplock bag
- 5 B. Allyl Ester Deprotection/1st Amide Formation:
1. Swell resin (in each well) with CHCl₃ (3x0.5 mL, 3 min); drain
 2. Add CHCl₃ solution of N-methylaniline (0.5 M, 0.5 mL)
 3. Add CHCl₃ solution of Pd(PPh₃)₄ (0.05 M, 0.5 mL)
 4. Bubble N₂ & vortex for 45 min; drain
 - 10 5. Wash with CHCl₃ (3x0.5 mL, 3 min)
 6. Repeat steps 2 – 5
 7. Wash resin with DMF solution of diethyldithiocarbamic acid, sodium salt trihydrate (0.03 M) and DIEA (0.06 M) (3x0.75 mL, 3 min)
 8. Wash resin with DMF (3x0.5 mL, 3 min)
 - 15 9. Wash resin with NMP (3x0.5 mL, 3 min)
 10. Add the following: NMP solution of DIEA (1.25 M, 0.15 mL), NMP solution of HATU (0.5 M, 0.15 mL) and NMP solution of amine (0.5 M, 0.15 mL) respectively (amine (HCl)_z were treated with an excess of an NMP solution of DIEA (1.25 M, (z)x0.15 mL)
 11. Bubble N₂ & vortex for 2 h; drain
 - 20 12. Wash with NMP (3x0.5 mL, 3 min)
 13. Repeat steps 10 – 12
 14. Wash resin with 1:1 NMP/dichloromethane (2x0.5 mL, 3 min)
 15. Wash resin with neat dichloromethane (3x0.5 mL, 3 min)
 16. Keep 96-well plate in the reaction block at rt overnight
- 25 C. TMSE Ester Deprotection/2nd Amide Formation:
1. Swell resin with THF (3x0.5 mL, 3 min); drain
 2. Add THF solution of TBAF (1 M, 0.5 mL)

3. Bubble N₂ & vortex for 45 min; drain
4. Wash resin with THF (3x0.5 mL, 3 min)
5. Repeat steps 2 – 4
6. Wash with 1:1 THF/NMP (2x0.5 mL, 3 min)
- 5 7. Wash with NMP (3x0.5 mL, 3 min)
8. Add the following: NMP solution of DIEA (1.25 M, 0.15 mL), NMP solution of HATU (0.5 M, 0.15 mL) and NMP solution of amine (0.5 M, 0.15 mL) respectively (amine (HCl)_z were treated with an excess of an NMP solution of DIEA (1.25 M, (z)x0.15 mL))
9. Bubble N₂ & vortex for 1.5 h; drain
- 10 10. Wash with NMP (3x0.5 mL, 3 min)
11. Repeat steps 8 - 10
12. Wash with 1:1 NMP/dichloromethane (2x0.5 mL, 3 min)
13. Wash with neat dichloromethane (3x0.5 mL, 3 min)
14. Store at about -15/-20°C in a ziplock bag
- 15 D. TFA Cleavage of Compound from Resin:
 1. Swell resin with dichloromethane (2x0.5 mL, 2 min); drain
 2. Add 95:5 TFA/H₂O solution to each well (0.5 mL)
 3. Bubble N₂ & vortex for 2 h; drain into cube tubes
 4. Wash wells with TFA/H₂O (3x0.25 mL, 2 min)
 - 20 5. Add AcOH (0.5 mL) to each cube tube
 6. Concentrate under reduced pressure with heat (Savant) for about 1 h
 7. Add AcOH (0.75 mL) to each cube tube
 8. Concentrate under reduced pressure with heat (Savant) for 45 min
 9. Add AcOH (0.25 mL) and toluene (0.75 mL) to each cube tube
 - 25 10. Concentrate under reduced pressure with heat (Savant) for 2 h
 11. Add methanol (0.25 mL), vortex then add toluene (0.75 mL)
 12. Concentrate under reduced pressure with heat (Savant) overnight

A 1152-member bicyclic library was produced using TentaGel™ as the solid support and the procedure described in steps A-D above. The library was made using 36 (3*12) by 32 (4*8) sets of diverse amines (see Table 2). The acid-labile protecting groups tert-butoxycarbonyl, tert-butyl ethers, and tert-butyl esters were utilized for the protection of amines, alcohols and carboxylic acids, respectively. On average, each well provided 6.5 micromoles of desired product [17.2 micromoles (of starting resin) * 0.76 (% yield) * 0.5 (assuming 50% purity on average)]. Each well was analyzed by MS (loop injection). In addition, 15 wells from plate 4 and 12 wells from plate 12 were analyzed by LC-MS to confirm that MS-loop injection analysis was consistent with the LC-MS data. Each compound of the 1152-member library was then placed into one of three relative purity categories: high purity, lower purity and failures. The data is summarized in Table 1.

Table 1


Plate #	# of High Purity wells	%	# of Lower Purity wells	%	# of Failures	%
1	69	72	18	19	9	9
2	77	79	18	19	1	2
3	74	77	22	23	0	0
4	74	77	11	11	11	11
5	72	75	21	22	3	3
6	68	71	28	29	0	0
7	76	79	20	21	0	0
8	77	80	19	20	0	0
9	78	81	18	19	0	0
10	67	70	29	30	0	0
11	49	51	47	49	0	0
12	62	65	34	35	0	0
Totals	843	73	285	25	24	2










High purity indicates that the molecular ion and/or fragments resulting from the desired ion were the only/major peaks in the MS spectra. Lower purity refers to wells where the molecular ion and/or fragment were present in addition to a number of other










peaks. Although a significant number of wells were of lower purity, the major impurity in these wells (about 90% of the wells) was the carboxylic acid resulting from incomplete coupling with the second amine. A failure indicates very little or no molecular ion or identifiable fragment was detected.















Table 2















Structures of Combinatorial Library Compounds



	NRR'
1	
2	
3	
4	
5	
6	$N(C_5H_{11})_2$
7	
8	
9	
10	

	NR''R'''
1	
2	$N(C_5H_{11})_2$
3	
4	
5	
6	
7	
8	
9	
10	

11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	

11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	

11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	

11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	

EXAMPLE 107

BIOLOGICAL ACTIVITIES OF REPRESENTATIVE BICYCLOOCTANES

Apoptosis:

The protocol used for determining inhibition of apoptosis in A549 cells was adopted from a system previously described (K. Last-Barney *et al.*, *J. of Immunology* 141:527-530, 1988). Briefly, 10^5 cells in 200 μ L 10% FBS/RPMI antibiotic containing culture medium were plated into 96 well round bottom culture plates and allowed to adhere for 6 hours at 37°C in a 5% CO₂ atmosphere. The media was removed and 100 μ L of RPMI + 1 μ g/mL actinomycin-D was added to each well, followed by 90 μ L of test compound solution in 1% DMSO. This was incubated for 1 hour. 10 μ L TNF- α was added at its EC₅₀ (normally 1 ng/mL FAC) and the plates incubated for 18 hours. The media was aspirated from the plates and 100 μ L of 0.5% crystal violet in 20% methanol was added. After 10 minutes the plates were rinsed with water to remove excess stain, air dried, and read on a Spectramax at a wavelength of 590 nm. The data obtained from the Spectramax was converted into percent inhibition data at a concentration of 20 μ M or IC₅₀ measured in μ M. Data is presented for representative compounds under the column titled "Apopt inh" in Table 3 as follows: "*" refers to percent inhibition from 6% to 64%; "***" refers to an IC₅₀ from 10 μ M to 50 μ M; "****" refers to an IC₅₀ below 10 μ M.



NF κ B:

A549 cells were stably transfected with an E-selectin promoter containing three NF κ B binding sites driving luciferase expression. For the assay, 5×10^4 cells were incubated in 96 well round bottom plates overnight in 100 μ L of 10% FBS/RPMI medium at 37°C in a 5% CO₂ atmosphere. The following morning the medium was removed and 90 μ L of a 1% DMSO solution of test compound solution was added and the plates incubated for 1 hour. 10 μ L of TNF- α was added at its EC₅₀ (normally 6 ng/mL FAC) to each well and the plate incubated for 5 hours. 100 μ L of luciferase buffer was added, and







after 10 minutes luminescence was read on a Wallac Victor 1420 Multilabel Counter. The data obtained from the Wallac Victor was converted into % inhibition data or IC₅₀ measured in μ M. Data is presented for representative compounds under the column titled "NF κ B inh" in Table 3 as follows: "*" refers to percent inhibition from 6% to 64%; "***" refers to an IC₅₀ from 10 μ M to 50 μ M; "****" refers to an IC₅₀ below 10 μ M.






The compounds of Table 3 were synthesized according to disclosed methods of Examples 1-106, and tested for activity according to the above assays. In Table 3, each compound is provided with a unique compound number, as set forth in the column "No."






Table 3







STRUCTURE	No.	Apopt inh	NF κ B inh
	1	**	*
	2	*	*







STRUCTURE	No.	Apopt inh	NFκB inh
	3	*	*
	4	*	*
	5	**	*
	6	*	*






STRUCTURE	No.	Apopt inh	NFκB inh
	7	*	*
	8	*	**
	9	*	*
	10	***	**
	11	**	*
	12	**	**







STRUCTURE	No.	Apopt inh	NFκB inh
	13	**	*
	14	**	**
	15	**	*
	16	**	**
	17	*	*







STRUCTURE	No.	Apopt inh	NFκB inh
	18	**	—
	19	*	*
	20	*	*
	21	*	*
	22	—	*






STRUCTURE	No.	Apopt inh	NFκB inh
	23	—	**
	24	—	**
	25	**	*
	26	—	*
	27	**	*
	28	**	*






STRUCTURE	No.	Apopt inh	NFκB inh
	29	**	*
	30	***	—
	31	*	*
	32	**	*
	33	**	*
	34	***	**






STRUCTURE	No.	Apopt inh	NFκB inh
	35	***	*
	36	*	*
	37	**	*
	38	***	**
	39	*	**







STRUCTURE	No.	Apopt inh	NFκB inh
	40	**	*
	41	*	**
	42	*	*
	43	*	*
	44	*	*
	45	***	*





STRUCTURE	No.	Apopt inh	NFκB inh
	46	*	*
	47	*	*
	48	**	*
	49	*	*
	50	***	**
	51	***	**

STRUCTURE	No.	Apopt inh	NFκB inh
	52	*	**
	53	***	**
	54	***	**
	55	—	*
	56	***	*






STRUCTURE	No.	Apopt inh	NFκB inh
	57	**	*
	58	**	*
	59	***	**
	60	**	**
	61	**	*






STRUCTURE	No.	Apopt inh	NFκB inh
	62	**	*
	63	*	*
	64	***	*
	65	***	**
	66	*	**






STRUCTURE	No.	Apopt inh	NFκB inh
	67	***	**
	68	***	**
	69	***	*
	70	*	*
	71	***	**
	72	***	*

STRUCTURE	No.	Apopt inh	NFκB inh
	73	—	**
	74	—	**
	75	—	*
	76	—	**







STRUCTURE	No.	Apopt inh	NFκB inh
	77	—	**
	78	***	—
	79	***	**
	80	***	**
	81	***	*




STRUCTURE	No.	Apopt inh	NFκB inh
	82	***	**
	83	***	**
	84	***	**
	85	—	**
	86	—	**

STRUCTURE	No.	Apopt inh	NFκB inh
	87	—	**
	88	—	*
	89	—	*
	90	—	*
	91	—	**

STRUCTURE	No.	Apopt inh	NFκB inh
	92	—	**
	93	—	**
	94	—	**
	95	*	*
	96	***	*

STRUCTURE	No.	Apopt inh	NFκB inh
	97	*	**
	98	*	*
	99	—	**
	100	—	**
	101	***	*

STRUCTURE	No.	Apopt inh	NFκB inh
	102	*	*
	103	*	*
	104	—	*
	105	—	*
	106	—	**
	107	—	**

STRUCTURE	No.	Apopt inh	NFκB inh
	108	—	**
	109	—	**
	110	*	*

EXAMPLE 108

BIOLOGICAL ACTIVITIES OF REPRESENTATIVE BICYCLOOCTANES

Compounds of the present invention were synthesized according to methods disclosed in Examples 1-106, and tested for activity according to the apoptosis and NFκB assays described in Example 107, and the CXCR1 and CXCR2 assays described below. The results from these biological testings are set forth in Table 4, where each compound is provided with a unique compound number, as set forth in the column “No.”


CXCR1:




This assay is a radioligand binding assay in human recombinant CHO cells with ^{125}I labeled IL-8 as the ligand. The assay procedure is described in Ahuja, S.K.; Murphy, P.M.; *J. Biol. Chem.* 1996, **271**, 20545, and was performed by Panlabs Taiwan, Ltd. Data is presented for representative compounds under the column titled "CXCR1 inh" in Table 4 as follows: "*" refers to percent inhibition from 10-36%; "**" refers to an IC_{50} from 10 μM to 50 μM ; "***" refers to an IC_{50} below 10 μM .



CXCR2:

This assay is a radioligand binding assay in human recombinant CHO cells with ^{125}I labeled IL-8 as the ligand. The assay procedure is described in Ahuja, S.K.; Murphy, P.M.; *J. Biol. Chem.* 1996, **271**, 20545, and was performed by Panlabs Taiwan, Ltd. Data is presented for representative compounds under the column titled "CXCR2 inh" in Table 4 as follows: "*" refers to percent inhibition from 10-36%; "**" refers to an IC_{50} from 10 μM to 50 μM ; "***" refers to an IC_{50} below 10 μM .

Table 4

STRUCTURE	No.	Apopt. Inh.	NFkB inh.	CXCR1 inh.	CXCR 2 inh.
	111	***	**	***	***

STRUCTURE	No.	Apopt. Inh.	NFkB. inh.	CXCR1 inh.	CXCR 2 inh.
	112	N.D.	*	**	***
	113	N.D.	*	**	**
	114	N.D.	*	N.D.	*

STRUCTURE	No.	Apopt. Inh.	NFkB inh.	CXCR1 inh.	CXCR 2 inh.
	115	—	*	*	*
	116	***	**	*	*

All other acronyms and abbreviations have the corresponding meaning as published in journals relative to the art of organic chemistry. From the foregoing it will be appreciated that, although specific embodiments of the invention have been described

 5 herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

All references cited herein are incorporated by reference.